

Genetic Diversity of Some Isolates of Soft rot *Pectobacterium*, from Different Host Plants in Fars Province, Iran, using RAPD Marker

Negin Akramipour¹, Mohammad Abdollahi^{1*},
S. Mohsen Taghavi² and Rasoul Rezaei¹

¹Department of Plant Protection, College of Agriculture, Yasouj University, Yasouj, Iran.

²Department of Plant Protection, College of Agriculture, Shiraz University, Shiraz, Iran.

(Received: 11 April 2016; accepted: 24 May 2016)

Pectobacteria are one of the most important groups of plant pathogenic bacteria which occur world-wide. The bacteria have a wide host range and infect plants with fleshy organs. Thirty five bacterial isolates were collected from potato, cabbage, sugar beet, pepper, carrot, onion, cucumber, turnip, and tomato host plants cultivated in Fars province. The bacteria were identified using biochemical and physiological tests. The genetic diversity among the isolates was evaluated using RAPD assay. The isolates were rod-shape, gram-negative, facultative anaerobes, oxidase-negative, catalase-positive, and had ability to rot the potato, while they could not induce hypersensitive reaction on tobacco leaf. All of isolates were identified as soft-rot bacteria. Based on phenotypic criteria, the isolated strains were categorized into three groups. The strains of first, second and third groups were related to *Pectobacterium carotovorum* subsp. *carotovorum*, *Pectobacterium betavascularum* and *Dickeya chrysanthemi*, respectively. According to RAPD assay and using six primers including 19300, 91200, OPB07, OPB08, OPB11, OPB18, isolates categorized in three groups. The used primers showed significant diversity among the strains and the specific primers GTGC4, GAGC4 and GTG5 confirmed the classification. The diversity among the isolates has been confirmed based on the molecular works and clustering based on RAPD. This is the first RAPD assay to investigate the genetic diversity of soft-rot bacteria.

Keywords: *Dickeya*; genetic diversity; pectobacteria; RAPD; soft-rot.

Pectobacteria, the causal agents of soft-rot, belonging to *Enterobacteriaceae* and cause many losses in most of crops, ornamental plants and vegetables (Alpini *et al.*, 1997; Charkowski, 2006; De Haan *et al.*, 2008; Duarte *et al.*, 2004; Kim *et al.*, 2009; Pitman *et al.*, 2008). They are able to produce cell wall degrading enzymes, especially pectolytic enzymes and can cause decay of parenchyma tissues (Permabelon & Salmond, 1995). Bacterial soft-rot occurs on plants with soft and fleshy storage parts or leaves and stems. Symptoms in diseased plants are very similar in all types of hosts. The bacterium can cause various damages

including reduce in germination, chlorosis, necrosis, stem rot, blackleg and eventually plant death (Avrova *et al.*, 2002; Helias *et al.*, 2000). Based on molecular and biochemical tests and host range, the genus *Pectobacterium* contains some species and subspecies including *P. carotovorum* subsp. *carotovorum*, *P. carotovorum* subsp. *odoriferum*, *P. brasiliensis*, *P. betavascularum*, *P. atrosepticum* and *P. wasabiae* (Garden *et al.*, 2003; Ma *et al.*, 2007; Van Der Merwe *et al.*, 2010). In soft-rot pectobacteria, two species, *Pectobacterium carotovorum* subsp. *carotovorum* and *Dickeya chrysanthemi*, have more economic importance and wider host rang. These two species infect crops of tropical and subtropical regions while *P. atrosepticum* exclusively is found associated with stem of

* To whom all correspondence should be addressed.
E-mail: rrezaei@yu.ac.ir

potatoes and cause blackleg disease in temperate climate (Permobelon, 1992). Bacterial soft-rot disease first reported from carrot by Jones in 1901 in America. In 1902, Van Hall from Netherland and Apple from Germany independently introduced the blackleg disease from potato (Goto & Motsumoto, 1987; Graham, 1964). In Iran, *Erwinia* was first isolated from rotted corms of cyclamen (Hedjarood, 1967). By phenotypic tests, the isolated bacterium was identified as *E. carotovora* (Amani, 1969). Some other studies were conducted to identify the isolates from different region of Iran. Most of these investigations were done based on physiological and biochemical tests, comparisons of the patterns of protein electrophoresis and the length of PCR amplified fragment using specific primers (Bahar & Danesh, 1986; Soltaninejad *et al.*, 2005; Zohoor *et al.*, 1998).

Performing molecular and biochemical tests and also host preference, is of importance to achieve accurate taxonomy, diagnosis, pathogen detection and epidemiological studies (Van Der Merwe *et al.*, 2010). Many studies have been conducted for taxonomical studies and identification of the soft rot bacteria so far. Some molecular methods such as RFLP (Darrasse *et al.*, 1996; Sleds *et al.*, 2000), AFLP (Avrova *et al.*, 2002), rep-PCR (Toth *et al.*, 1999), RAPD (Maki Valkama & Karjalainen, 1994) and the analysis of 16S rRNA sequences (Hauben *et al.*, 1998) have been done to characterization of soft rot bacteria.

This study was aimed to use phenotypic characteristics as well as RAPD marker to evaluate the genetic diversity among the soft-rot bacteria isolated from potato, cabbage, carrot, pepper, tomato, cucumber, turnip and date palm tree in Fars province.

MATERIALS AND METHODS

Sampling

During the summer months of 2013, the samples were collected from potato, cabbage, carrot, pepper, tomato, cucumber, turnip and date palm trees with the symptoms of pectobacterial infection, in different regions of Fars province. Two reference isolates (EccSCR and EchSCR) which were provided by Scottish Crop Research Institute were used to compare with the isolated bacteria. The information on the isolates is given in Table 1.

Isolation and purification

In order to isolate the bacteria, the infected plant parts were washed with tap water. Bacteria were isolated from the interface between healthy and infected portions of the rotted tissue. The tissues were dipped in 10% sodium hypochlorite for 30 s and then washed with sterile distilled water to disperse the bacterial cells into water. Finally, the pieces were transferred into sterilized Petri dishes containing a few ml of distilled water and was shaken for 15-30 min. The content of Petri dishes were transferred into sterilized mortar and then homogenized. This suspension is then streaked on nutrient agar (NA) medium and the Petri dishes were incubated at 27-30°C. After 24-48 h, to purify the isolated bacteria, the relatively slimy and milky small colonies were sub-cultured on fresh NA medium. The small and relatively embossed colonies with bright green color were transferred to Eosin Methylene-Blue (EMB) medium and the purified colonies were obtained.

Investigation of phenotypic characteristics

To identify the soft-rot pectobacteria, the tests of gram, oxidase, urease, catalase, potato rot, hypersensitive reaction and growth at 37 and 40°C, reductive materials production from sucrose and indole and phosphate production were conducted according to Schaad *et al.* (2001), and aerobic/anaerobic growth assay was carried out as described by Hugh & Leifson (1953). Erythromycin sensitivity assay was carried out based on the method described by Kelment *et al.* (1964).

DNA extraction

DNA was extracted by method described by Kaung *et al.* (2003). Bacteria were grown on LB medium (0.5% yeast extract, 1% Trypton, 1% Sodium Chloride), for 20-24 h. Three ml of bacterial culture was centrifuged at 14000 rpm for three min. The pellet was dissolved in 500 µl of CTAB buffer (2 g of CTAB, 0.6 g of Tris, 2.1 g of NaCl and 0.4 g of EDTA) and heated at 65°C for 30 min. Chloroform-isoamylalcohol solution (1:24) was added into the tubes and centrifuged at 13000 rpm for eight min, then the supernatants were transferred into new tubes. Equal volumes of cold isopropanol were added into the tubes and centrifuged at 13000 rpm for eight min for DNA precipitation. Finally, the tubes were washed using 70% ethanol and then were dried. 50 µl of TE buffer (Tris-EDTA) or

sterilized still water were added to the tubes and stored at -20°C. Two µl of the extracted DNA was subjected to PCR.

Pathogen identification by using specific primers

Identifying the bacteria was carried out using Y1/Y2 specific primers (Y1: 5' TTACCGGACGC CGAGCTGTGGCGT 3'/Y2: 5' CAGGAAGATGTCGTTATCGCGAGT 3') (DARRASSE *et al.* 1996), with annealing temperature of 65°C and expected product size of 434 bp. *P. carotovorum* subsp. *carotovorum* was also identified using EXPCCR/EXPCCF specific primers (EXPCCR: 5' GCCGTAATTGCC TACCTGCTTAAG 3'/ EXPCCF: 5' GAACTT CGCACCGCCGACCTTCTA 3') (Kaung *et al.*, 2003), with annealing temperature of 60°C and expected product size of 550 bp. Finally, ADE1/ADE2 specific primers (ADE1:5' GATCAGA AAGCCCGCAGCCAGAT/3'/ADE2:5'CTGTGGC CGATCAGGATGG 3') with annealing temperature of 65°C and expected product size of 434 bp were used to identify *Dickeya* spp. (Nasser *et al.*, 1994). The primers were generated in Pishgam Biotech Company, Iran. PCR was performed in a reaction mixture of 25 µl, containing 2 µl of DNA, 2.5 µl of 10X PCR Buffer, 1.5 µl of MgCl₂, 0.7 µl of dNTPs (25 mM), 0.5 µl of *Taq* DNA Polymerase and 1 µl of each primer (10 pmol). PCR products were stored at -20°C until gel electrophoresis.

Study on genetic diversity using RAPD

To compare the genetic diversity in the populations, RAPD analysis carried out by using six short random primers (Table 2). The assay was performed in a reaction mixture of 25 µl, containing 2 µl of DNA, 2.5 µl of 10X PCR Buffer, 2.5 µl of MgCl₂, 0.7 µl of dNTPs (25 mM), 0.5 µl of *Taq* DNA Polymerase and 1 µl of each primer (10 pmol). PCR was performed in a thermocycler for all primer. One cycle of initial denaturation at 94°C for one min, 40 cycles including denaturation at 94°C for one min, annealing at 35°C for one min and extension at 72°C for two min, and a final extension at 72°C for 10 min. PCR products were stored at -20°C until gel electrophoresis.

Gel electrophoresis

To carry out PCR reaction, 6-7 µl of each PCR product were mixed with 2 µl of loading buffer and were electrophoresed in 1% agarose gel at 80-90 volt for 1 h. Ruler 100 bp DNA ladder was used for determination of the fragments size. The gel

was dyed using ethidium bromide and then destained in still water. The stained gels were observed with the UV 254 transilluminator and photographed.

Analysis of molecular data

The genetic similarities among strains were analyzed by NTSYS-PC ver. 2.02 software (Numerical Taxonomy and Multivariate Analysis System), on the basis of electrophoresis profiles. Genetic distance or similarity among the strains based on molecular markers, was scored as the presence/absence of band in the gel. The matrix of similarity of coefficients was subjected to un-weighted pair-group method analysis (UPGMA) to generate dendrogram, using the average linkage procedure. (Rohlf, 2000).

RESULTS

Phenotypic characteristics

All isolates were rod-shape, gram-negative, facultative anaerobe, oxidase-negative, catalase-positive and were able to cause potato rot while they could not induce hypersensitive reaction. They did not produce yellow colonies on YDC medium, but they showed bright green colonies on EMB medium (Gallios *et al.*, 1992; Schaad *et al.*, 2001). According to differential physiological characteristics (Table 3), the isolates were categorized into three groups. The first group which grown at 37°C but not at 40°C and were negative for indole production, sucrose reduction, phosphatase activity and insensitive to erythromycin, were similar to *P. carotovorum* subsp. *carotovorum* (*Pcc*) (Fahy & Persly, 1983). The second group were grown at 37°C, unable to grow at 40°C and were negative for indole production, phosphatase activity and insensitive to erythromycin, but positive to sucrose reduction, were similar to *Pectobacterium betavasculorum* (*Pb*) (Thomson & Schroth, 1981). Third group were grown at 40°C, were positive for indole production, phosphatase activity and sensitive to erythromycin, were found to be similar to *Dickeya chrysanthemi* (*Dch*) (Nasser *et al.*, 1994).

PCR reaction and characterization of the bacterial isolates

Except for *P. betavasculorum*, the primers Y₁/Y₂ were amplified the expected product size of 434 bp, and characterized the *Pectobacterium*

Table 1. *Pectobacterium* isolates from different hosts in Fars Province

Isolate	Origin	Host	Isolate	Origin	Host
O ₃	Marvdasht	Onion	S ₂	Zargan	Sugar beet
Tu ₁	Kazerun	Turnip	S ₃	Zargan	Sugar beet
Tu ₂	Shiraz	Turnip	S ₄	Zargan	Sugar beet
Tu ₃	Kazerun	Turnip	S ₅	Zargan	Sugar beet
P ₅	Marvdasht	Potato	S ₆	Zargan	Sugar beet
P ₆	Marvdasht	Potato	P ₁	Kazerun	Potato
Cu ₂	Kazerun	Cucumber	P ₂	Kazerun	Potato
C ₃	Shiraz	Carrot	Cu ₁	Marvdasht	Cucumber
C ₄	Fasa	Carrot	Pa ₁	Farashband	Palm
Pe ₁	Fasa	Pepper	O ₁	Fasa	Onion
Pe ₂	Fasa	Pepper	Pa ₂	Farashband	Palm
Ca ₁	Marvdasht	Cabbage	O ₂	Farashband	Onion
Ca ₂	Marvdasht	Cabbage	P ₇	Shiraz	Potato
To ₁	Shiraz	Tomato	P ₈	Shiraz	Potato
To ₂	Marvdasht	Tomato	C ₁	Mianrud	Carrot
Pa ₄	Asaluyeh	Palm	C ₂	Mianrud	Carrot
EccSCR	Scotland	Potato	P ₃	Kaftarak	Potato
EchSCR	Scotland	Potato	P ₄	Kaftarak	Potato
S ₁	Shiraz	Sugar beet			

isolates (Fig. 1). Moreover, by amplifying the expected 550 bp products, the oligonucleotide primers EXPCCR/EXPCCF found to be suitable for separating *Pcc* that isolated from various hosts (Fig. 2).

RAPD assay

RAPD assay was performed using the primers 19300, 91200, OPB07, OPB08, OPB11, OPB18. All primers successfully amplified DNA fragments and clearly showed genetic diversity among soft-rot isolates (Figs. 3-5). Complementary analysis using genetic fingerprinting from three RAPD primers showed that the isolates fall into three groups (Figs. 6 and 7). The first group includes all *Pb* isolates which produced exclusive electrophoresis patterns. The second and the third

groups include *Dch* and all of *Pcc* isolates, respectively. In this way, thirty five isolates from various hosts which were identified as *Pcc*, *Pb* and *Dch* using phenotypic characteristics, were easily identified by RAPD assay.

Table 2. Primers used in RAPD test for characterization of *Pectobacterium* isolates from different hosts in Fars Province

Primer name	Primer sequence
19300	CGATTCGGCG
91200	GATAACGCAC
OPB-07	GGTGACGCAG
OPB-08	GTGACGTAGG
OPB-11	GTAGACCCGT
OPB-18	CCACAGCAGT


**Fig. 1.** DNA fingerprinting pattern of *Pectobacterium* sp. isolates generated by Y₁ and Y₂ primers 1: Water control, 2: Standard isolate, 3: P₁, 4: Pa₁, 5: C₁, 6: P₃, 7: O₃, 8: Tu₁, 9: P₅, 10: Pe₁, 11: To₁, 12: C₃, 13: Ca₁, M: Marker (100bp)

Table 3. Differential physiological properties of *Pectobacterium* isolates from different hosts in Fars Province

Test	Bacterial isolates																				
	S ₁	S ₂	S ₃	S ₄	S ₅	S ₆	S ₇	S ₈	S ₉	S ₁₀	S ₁₁	S ₁₂	S ₁₃	S ₁₄	S ₁₅	S ₁₆	S ₁₇	S ₁₈	S ₁₉	S ₂₀	
Gram test	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Oxidase production	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Catalase production	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Aerobe growth	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Anaerobe growth	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Urease production	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Potato soft rot	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
HR on Tobacco	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Growth at 37°C	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Growth at 40°C	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Yellow pigment on YDC	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Phosphatase activity	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Indole production	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Erythromycin sensitivity	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Reducing substrate from sucrose+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Pectobacterium</i> strain	Pb	Pb	Pb	Pb	Pb	Pb	Pb	Pb	Pb	Pb	Pb	Pb	Pb	Pb	Pb	Pb	Pb	Pb	Pb	Pb	Pb
<i>Pectobacterium</i> betavasculorum																					
<i>Dickeya chrysanthemi</i>																					
<i>Pectobacterium carotovorum</i> subsp. <i>carotovorum</i>																					




Fig. 2. DNA fingerprinting pattern of *Pectobacterium carotovorum* subsp. *carotovorum* isolates generated by EXPCCR and EXPCCF primers M: Marker (100bp), 1: Water Control, 2: P₁, 3: P₂, 4: Pa₁, 5: O₁, 6: P₃, 7: O₃, 8: C₁, 9: C₅, 10: P₄, 11: To₁, 12: Standard isolate




Fig. 3. DNA fingerprinting pattern of isolates generated by RAPD and primer 193001: S₁, 2: S₂, 3: S₃, 4: S₄, 5: S₅, 6: S₆, 7: EchSCR, 8: O₂, 9: Pa₂, 10: P₇, 11: P₈, 12: P₁, 13: P₂, 14: Cu₂, 15: O₁, 16: Pa₁, M: Marker (100bp).




Fig. 4. DNA fingerprinting pattern of isolates generated by RAPD and primer OPB07 M: Marker (100 bp), 1: C₁, 2: C₂, 3: P₃, 4: P₄, 5: O₃, 6: Tu₁, 7: Tu₂, 8: Tu₃, 9: Cu₂, 10: P₅, 11: P₆, 12: C₃, 13: C₄, 14: Pe₁, 15: Pe₂, 16: Ca₁, 17: Ca₂, 18: To₁, 19: To₂, 20: Pa₄.




Fig. 5. DNA fingerprinting pattern of isolates generated by RAPD and primer OPB08
 1: S₁, 2: S₂, 3: S₃, 4: S₄, 5: S₅, 6: S₆, 7: EchSCR, 8: O₂, 9: Pa₂, 10: P₇, 11: P₈, 12: Cu₁, 13: P₁, 14: P₂, 15: O₁, M: Marker (100bp)




Fig 6. DNA fingerprinting pattern of isolates generated by RAPD and primer OPB18
 1: C₁, 2: C₂, 3: P₃, 4: P₄, 5: O₃, 6: Tu₁, 7: Tu₂, 8: Tu₃, 9: P₅, 10: P₆, 11: Cu₂, 12: C₃, 13: C₄, 14: Pe₁, 15: Pe₂, 16: Ca₁, 17: Ca₂, 18: To₁, 19: To₂, 20: Pa₄, M: Marker (100bp)




Fig. 7. Dendrogram of soft rot *Pectobacterium* strains based on 6 primers of RAPD
 Pb: *Pectobacterium betavasculorum* Dch: *Dickeya chrysanthemi* Pcc: *Pectobacterium carotovorum* subsp. *carotovorum*

DISCUSSION

In this study, significant diversity was observed among the tested isolates, in case of phenotypic characteristics. However, in some cases, an inconsistency with previously identified species was found. It has reported earlier that bacterial identification seems to require more reliable, precise and faster methods, such as DNA-based techniques. According to Toth *et al.* (2001), RAPD method can be used as a faster and more suitable method than the phenotypic methods.

Our study showed that investigation of genetic diversity using RAPD assay is a reliable method for identification and classification of *Pectobacterium* spp. In the present research, the results obtained from the primers 19300, 91200, OPB07, OPB08, OPB11 and OPB18, showed that all of the primers successfully amplified the DNA fragments. Of these primers, 19300, 91200, OPB07 and OPB11 was able to differentiate the three species of the soft rot bacteria. Analysis of genetic fingerprinting from six RAPD primers showed that the isolates which have 66% similarity, are categorized into two groups. The first group is divided to two sub-group, first includes *Pb* isolates with 95% and *Dch* isolates with 85% similarity and the second includes some isolates of *Pcc* with 70% similarity. The other isolates of *Pcc* were fallen into the second group with 69% similarity. In our study, no PCR product was amplified in case of *Pb* and *Dch* isolates. This is agree with previous findings (Helias *et al.*, 2000; Kaung *et al.*, 2003).

RAPD assay of the *Pcc* isolates showed that these are genetically diverse isolates whereas the *Pb* isolates have more genetic homogeneity. More restricted host range, smaller population size and even similar origin may be the reasons for lesser genetic diversity of the *Pb* isolates.

In *Pcc* group, the isolates P₁, P₂, C₁ and C₂ which were isolated from a same host and region share 95% similarity. However, in some cases, it is possible that two isolates from a same host and a same geographical region fall into different groups with less similarity, indicating high level of diversity in a single geographical region.

Maki-Valkama & Karjalainen (1994) successfully used RAPD to distinguish *Erwinia carotovora* subsp. *carotovora* (*Ecc*) from *E. carotovora* subsp. *atroseptica* (*Eca*). They also

showed that the similarity among the isolates of *Eca* is more than 85% while high level of genetic diversity was observed among *Ecc* samples. Rezaei & Taghavi (2008) used rep-PCR with Eric, Box and rep primers and categorized the soft-rot pectobacteria into four groups: *Pcc*, *P. carotovorum* subsp. *odoriferum* (*Pco*), *Pb* and *Dch*. Furthermore, they showed that a high level of genetic diversity exists among *Pcc* isolates. Tavasoli *et al.* (2010) investigated the genetic diversity of the soft-rot pectobacterial population using BOX-PCR. Their results showed that there is a high level of diversity among the isolates.

Generally, the diversity has been found in genetic fingerprint of isolates from different regions. In contrast, isolates from a single region may fall into different groups. In a study, AFLP was performed by Avrova *et al.* (2002), and the tested pectobacteria were placed into three groups. The first includes *Pco* and *Pcc*, the second contains *P. atroseptica* (*Pa*) and *Pb* and the third harbors *Dch*. A polymerase chain reaction (PCR) test based on amplification of 16S–23S rRNA of intergenic transcribed spacer (ITS) characterization of *P. carotovorum* was carried out by Golkhandan *et al.* (2013). They results showed that PCR by G1/L1 followed by RCR-RFLP of G1/L1 primer using *RsaI* enzyme was a simple, rapid, and effective method for identification and differentiation of *Pcc* from *Dickeya* spp. and other species of *Pectobacterium*.

So far, RAPD assay has not been used to investigate the genetic diversity of soft-rot bacteria. However, this method has been used for some other plant pathogenic bacteria. Bysal *et al.* (2011) used ISSR method and showed that there is a high genetic diversity among the isolates of *Clavibacter michiganensis* subsp. *michiganensis* in Turkey. In another study, Samanta & Mandal (2013) were applied the ISSR test to show the genetic diversity among the isolates of *X. axonopodis* pv. *commiphorae*. Kawaguchi (2014) used ISSR to categorize the isolates of *X. arbuticola* pv. *pruni* into four groups, while the genetic diversity among the isolates was low.

The results of the present study showed that RAPD fingerprinting is a powerful tool to identify the various soft rot bacteria from different hosts and regions. The high genetic diversity was found in case of *Pcc* and *Dch* isolates. *Pb* isolates

were also found to have more genetic homogeneity.

ACKNOWLEDGMENTS

The research was supported by Research and Technology office and Post Graduate office of Yasouj University, Yasouj, Iran. The authors thank for their valuable assistance.

REFERENCES

- Alippi, A.M.E., Dal Bó, L., Ronco, B., Casanova, P.E. Tomato as a new host of *Erwinia carotovora* subsp. *carotovora* in Argentina, *Plant Disease*, 1997; **81**: 230.
- Amani, B. Rotting of ornamental plants and vegetables, *Iranian Journal of Plant Pathology*, 1969; **4**: 10-13. (In Farsi with English Summary).
- Avrova, A.O., Hyman, J., Toth, R.L., Toth, I.K. Application of amplified fragment length polymorphism fingerprinting for taxonomy and identification of the soft rot bacteria *Erwinia carotovora* and *Erwinia chrysanthemi*, *Applied and Environmental Microbiology*, 2002; **68**: 1499-1508.
- Bahar, M., Danesh, D. Occurrence of potato blackleg in Isfahan. Proceedings of 8th Iranian Plant Protec. Congress, 1986; Isfahan, Iran, pp 105. (Abst.).
- Charkowski, A.O. The Soft Rot *Erwinia*, In: S.S. Gnanamanickam (Ed.). Plant-associated bacteria. Springer, Dordrecht, The Netherlands, 2006; 423-505.
- Darrasse, A., Priou, S., Kotoujansky, A., Bertheau, Y. PCR and restriction fragment length polymorphism of a pel gene as a tool to identify *Erwinia carotovora* in relation to potato diseases, *Applied and Environmental Microbiology*, 1996; **60**: 1437-1443.
- De Haan, E.G., Dekker-Nooren, T.C.E.M., Van Den Bovenkamp, G.W., Speksnijder, A. G., Vander, C.L., Zouwen, P.S., Vander Wolf, J. M. *Pectobacterium carotovorum* subsp. *carotovorum* cause blackleg in temperate climate, *European Journal of Plant Pathology*, 2008; **122**: 561-569.
- Duarte, V., De Boer, S.H., Ward, L.J., Oliveira, A.M.R. Characteristic of atypical *Erwinia carotovora* strains causing black leg of potato in Brazil, *Journal of Applied and Microbiology*, 2004; **96**: 533-545.
- Fahy, P.C., Persly, G.J. Plant Bacterial Disease: A Diagnostic Guide. New York. Academic Press. 1983; 389pp.
- Fassihiani, A., Nedaeinia, R. Characteristics of Iranian *Pectobacterium Carotovora* strains from sugarbeet by phenotypic tests and whole-cell proteins profile. *Journal of Phytopathology*, 2008; **10**: 1-6.
- Gallios, A., Samson, R., Ageron, E., Grimont, P.A.D. *Erwinia carotovora* Subsp. *odorifera* subsp. nov., associated with odorous soft rot of chicory (*Cichorium intybus* L.), *International Journal of Systematic Bacteriology*, 1992; **42**: 582-588.
- Garden, L., Gouy, C., Christen, R., Samson, R. Elevation of three subspecies of *Pectobacterium betavascularum* sp. nov. and *Pectobacterium wasabiae* Sp. nov., *International Journal of Systematic Evolutionary and Microbiology*, 2003; **53**: 381-391.
- Golkhandan, E., Kamaruzaman, S., Serial, M. Characterization of Malaysian *Pectobacterium* spp. from vegetables using biochemical, molecular and phylogenetic methods. *European Journal of Plant Pathology*, 2013; **137**: 431-443.
- Goto, M., Matsumoto, K. *Erwinia carotovora* subsp. *wasabiae* subsp. nov. isolated from diseased rhizomes and fibrous roots of Japanese horseradish (*Eutrema wasabi* Maxim.). *International Journal of Systematic Bacteriology*, 1987; **37**: 130-135.
- Graham, D.C. Taxonomy of the soft rot coliform bacteria. *Annual Review of Phytopathology*, 1964; **2**: 13-42.
- Hauben, L., Moore, E.R.B., Vauterin, L., Steenackers, M., Mergaert, J., Verdonck, L., Swings, J. Phylogenetic position of phytopathogens within the Enterobacteriaceae. *Systemic Applied of Microbiology*, 1998; **21**: 384-397.
- Hedjarood, G. Soft rot of cyclamen in Iran. *Iranian Journal of Plant Pathology*, 1967; **4**: 19-23. (In Farsi with English Summary).
- Helias, V., Le Roux, A.C., Bertheau, Y., Andrivon, D.J., Gauthier, P., Jouan, B. Characterizations of *Erwinia carotovora* subspecies and detection of *Erwinia carotovora* subsp. *atroseptica* in potato plants, soil and water extracts with PCR-based methods. *European Journal of Plant Pathology*, 2000; **104**: 685-699.
- Hugh, R., Leifson, F. The taxonomic significance of fermentative versus oxidative metabolism of carbohydrates by various gram negative bacteria. *Journal of Bacteriology*, 1953; **66**: 24-26.
- Kaung, H.W., Kwon, S.W., Go, S.J. PCR-based specific and sensitive detection of *Pectobacterium carotovora* subsp. *carotovora* by primers generated from URP-PCR

- fingerprinting derived polymorphs band. *Plant Pathology*, 2003; **52**: 127-133.
21. Kelment, Z., Farkas, C., Lovrekorich, L. Hypersensitivity reaction induced by phytopathogenic bacteria in the tobacco leaf. *Phytopathology*, 1964; **54**: 474-477.
 22. Khademlou, E., Taghinasab, M., Taheri, A., Rahimian, H. Bacterial leaf spot of *Aglaonema* caused by strains resembling *Pectobacterium carotovorum* subsp. *odoriferum*. Proceedings of 17th Iranian Plant Protection Congress, Karaj, Iran. 2006; pp. 384. (In Farsi with English Summary).
 23. Kim, H.S., Ma, B., Perna, N.T., Charkowski, A.O. Prevalence and virulence of natural type III secretion system deficient *Pectobacterium* strains. *Applied Environmental Microbiology*, 2009; **75**: 4539-4549.
 24. Ma, B., Hibbing, M.E., Kim, H.S., Reedy, R.M., Yedidia, I., Breuer, J., Glasner, J.D., Perna, N.T., Kelman, A., Charkowski, A.O. Host range and molecular phylogenies of the soft rot enterobacterial genera *Pectobacterium* and *Dickeya*. *Phytopathology*, 2007; **97**: 1150-1163.
 25. Maki-Valkama, T., Karjalainen, R. Differentiation of *Erwinia carotovora* subsp. *atroseptica* and *carotovora* by RAPD-PCR. *Annals of Applied Biology*, 1994; **125**: 301-304.
 26. Nasser, Z., Bertheau, Y., Dervin, C.J., Narcy, P., Lemattre, M. Ribotyping of *Erwinia chrysanthemi* strains in relation to their pathogenic and geographic distribution. *Applied Environmental Microbiology*, 1994; **60**: 3781-3789.
 27. Perombelon, M.C.M. Potato blackleg: epidemiology, host-pathogen interaction and control. *Netherland Journal of Plant Pathology*, 1992; **98**: 135-146.
 28. Perombelon, M.C.M. Diversity in *Erwinias* as plant pathogens, In: M. Lemattre, S. Freigoun, K. Rudolph, and J.G. Swings. (Eds) Plant pathogenic bacteria. Versailles, France. Les Colloques 66, INRA-ORSTOM, Paris, 1994; 113-128.
 29. Perombelon, M.C.M., Salmond, G.P.C. Bacterial Soft Rots. In: U.S. Singh, R.P. Singh and K. Kohmoto, (Eds). Pathogenesis and host specificity in plant disease. Tottori University of Japan, 1995; 1-20
 30. Pitman, A.R., Wright, P.J., Galbraith, M.D., Harrow, S.A.C. Biochemical and genetic diversity of pectolytic enterobacteria causing soft rot disease of potatoes in New Zealand. *Australian Plant Pathology*, 2008; **37**: 559-568.
 31. Rezaei, R., Taghavi, S.M. Phenotypic and genotypic characteristics of Iranian soft rot bacteria isolates from different hosts. *Phytopathology Mediterranea*, 2010; **49** (2): 194-204.
 32. Rohlf, F.J. Ntsys-PC, numerical taxonomy and multivariate analysis system. version 2.11j. Setauket, NY. USA. 2000; Exeter Software.
 33. Samson, R., Legendre, J.B., Christen, R., Fischer-Le Saux, M., Achouak, W., Gardan, L. Transfer of *Pectobacterium chrysanthemi* (Burkholder *et al.*, 1953) Brenner *et al.*, 1973 and *Brenneria paradisiaca* to the genus *Dickeya* gen. nov. as *Dickeya chrysanthemi* comb. nov. and *Dickeya paradisiaca* comb. nov. and delineation of four novel species, *Dickeya Dadantii* sp. nov., *Dickeya dianthicola* sp. nov., *Dickeya dieffenbachiae* sp. nov. and *Dickeya zae* sp. nov. *International Journal of Systematic Evolutionary and Microbiology*, 2005; **55**: 1415-1427.
 34. Schaad, N. W., Jones, J.B., Chun, W. Laboratory Guide for Identification of Plant Pathogenic Bacteria. 3rd Edition. The American Phytopathological Society, St. Paul, Minnesota. USA., 2001; 373pp.
 35. Sleds, W., Jafra, S., Waleron, M., Lojkwaska, E. Genetic diversity of *Erwinia carotovora* strains isolated from infected plants grown in Poland. Bulletin, OEPP/EPPO Bulletin. 2000; **30**: 403-407.
 36. Soltaninejad, S., Taghavi, M., Hayati, J., Mostowfizadeh-Ghahamfarsa, R. Study of phenotypic features and pathogenicity of soft rot *Pectobacteria* in Khuzestan Province. *Iranian Journal of Plant Pathology*, 2005; **41**: 585-611. (In Farsi with English summary).
 37. Tavasoli, E., Marefat, A.R., Hasanzadeh, N. Phenotype and genotype diversity of soft rot *Pectobacteria* isolated from different plant hosts in several regions of Iran. *Biology Journal of Garmsar Azad University*, 2010; **4**(4): 69-78.
 38. Thomson, S. V., Schroth, M.N. Identification and differentiation of the *Erwinia* sugar beet pathogens from members of *Erwinia carotovora* and *Erwinia chrysanthemi*. *Phytopathology*, 1981; **71**: 1937-1042.
 39. Thomson, S.V., Schroth, M.N., Hills, F.J., Whitney, E.D., Hilbrand D.C. Bacterial vascular necrosis and rot of sugar beet: general description and etiology. *Phytopathology*, 1997; **67**: 1183-1189.
 40. Toth, I.K., Bertheau, Y., Hyman, L.J., Laplaze, L.M., López, M., McNicol, J., Niepold, F., Persson, P., Salmond, G. P. C., Sletten, A., Van Der Wolf, J. M., Pérombelon M. C. M. Evaluation of phenotypic and molecular typing techniques for determining diversity in *Erwinia*

- carotovora* subsp. *atroseptica*. *Journal of Applied Microbiology.*, 1999; **87**: 158-166
41. Van Der Merwe, J.J., Coutinho, T.A., Korsten, L., Van Der Waals, J.F. *Pectobacterium carotovorum* subsp. *brasiliensis* causing blackleg on potatoes. *European Journal of Plant Pathology.*, 2010; **126**: 175-185.
42. Zohoor, S.M., Rahimian, H., Banihashemi, N. Identity of soft rot *Erwinia* as infecting potato in Fars. Proceedings of 13th Iranian Plant Protection Congress, Karaj, Iran. 1998; 171p. (Abst.)