

Statistical Optimization of Phosphate Solubilization by *Erwinia* sp.hdds3fr

**Dalia El-Sayed El-Badan, Dunja-Manal Abou-Zeid,
Hind Mohamed Hassan and Soraya A. Sabry**

Botany and Microbiology Department, Faculty of Science, Alexandria University, Alexandria Egypt.

<https://doi.org/10.22207/JPAM.10.2.08>

(Received: 11 January 2015; accepted: 03 March 2015)

Phosphate solubilizing bacteria (PSB) are known to convert phosphate from the insoluble to soluble form, and make it available for plant uptake. The use of microbial inoculants (biofertilizers) possessing P-solubilizing activities in agricultural soils is considered as an environmental-friendly alternative to further applications of chemical based P fertilizers. Therefore, the objective of this research was to mine for phosphate solubilizing microorganisms from our local environments. Upon screening, 45 isolates were obtained showing varying degrees of phosphate solubilization on agar plates using PVK and NBIRP media. One bacterial isolate exhibited the highest phosphate solubilization on both media. The phosphorus solubilizing potential of the isolate was evaluated using solid and liquid media under in vitro conditions. The bacterial isolate was identified based on 16S rRNA gene sequencing data as *Erwinia* sp.hdds3fr. It was also aimed in this study to investigate nutritional factors affecting P solubilization using the Plackett-Burman experimental design. On the basis of the calculated t-values, the main factor that had significant positive effect on phosphate solubilization by *Erwinia* sp. hdds3fr was $\text{Ca}_3(\text{PO}_4)_2$. In order to validate the obtained data and to evaluate the accuracy of the applied Plackett-Burman statistical design, a verification experiment was carried out. A significant increase near 2 fold in SP was achieved.

Keywords: Phosphate solubilization; *Erwinia* sp. hdds3fr; optimization;
Plackett-Burman statistical design.

Phosphorus (P) is one of the major plant nutrients limiting plant growth, next to nitrogen (N). It plays an important role in virtually all major metabolic processes in plant including photosynthesis, energy transfer, signal transduction, macromolecular biosynthesis and respiration^{1,2} and nitrogen fixation in legumes³. Although P is abundant in soils in both inorganic and organic forms, its availability is restricted as it occurs mostly in insoluble forms. These insoluble, precipitated forms cannot be absorbed by plants⁴. The P content in average soil is about 0.05% (w/w)

but only 0.1% of the total P is available to plant because of poor solubility and its fixation in soil⁵. Moreover, the efficiency of applied P fertilizers in chemical form rarely exceeds 30% due to its fixation, either in the form of iron/aluminium phosphate in acidic soils⁶ or in the form of calcium phosphate in neutral to alkaline soils⁷. It has been suggested that the accumulated P in agricultural soils would be sufficient to sustain maximum crop yields worldwide for about 100 years if it were available^{8,9,10}. Furthermore P is a finite resource and based on its current rate of use, it has been estimated that the worlds known reserves of high quality rock P may be depleted within the current century¹¹. Beyond this time, the production of P based fertilizers will require the processing of lower grade rock at significantly higher cost¹². The realization

* To whom all correspondence should be addressed.
E-mail: daliaelbadan@hotmail.com

of all these potential problems associated with chemical P fertilizers together with the enormous cost involved in their manufacture, has led to the search for environmental compatible and economically feasible alternative strategies for improving crop production in low or P-deficient soils ¹³. The use of microbial inoculants (biofertilizers) possessing P-solubilizing activities in agricultural soils is considered as an environmental-friendly alternative to further applications of chemical based P fertilizers.

The objective of the present investigation was to isolate PSB that could solubilize insoluble phosphate efficiently and optimizing its culture conditions to improve solubilization process using Plackett-Burman design.

MATERIALS AND METHODS

Isolation of phosphate solubilizing bacteria

Soil employed to isolate PSM was collected from garden at Faculty of Science, Alexandria University in Egypt. One gm of soil was suspended in 9 ml distilled water and serially diluted up to 10^{-5} . A sample (100 μ l) from each serial dilution was plated on agar plates of PVK medium¹⁴ containing in gL⁻¹, glucose, 10; $\text{Ca}_3(\text{PO}_4)_2$, 5; $(\text{NH}_4)_2\text{SO}_4$, 0.5; NaCl, 0.2; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.1; KCl, 0.2; yeast extract, 0.5; $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, 0.002; $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.002; and on NBRIP medium¹⁵ containing in gL⁻¹ glucose, 10; tricalcium phosphate (TCP), 5; $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$, 5; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.25; KCl, 0.2; $(\text{NH}_4)_2\text{SO}_4$, 0.1. Medium pH was adjusted to 7 with HCl. Plates were incubated for 2 to 3 days at 30 °C. Colonies with clear halos were considered as phosphate solubilizing microorganisms (PSM)¹⁶. A total of 45 isolates that exhibited clear zones on the agar plates were selected. Based on the size of the clear zone, isolate H3fr was the most efficient phosphate solubilizer, and was selected for further studies.

Identification of isolate H3fr

Partial sequencing of 16S rRNA for the bacterial isolate H3fr was performed with the help of a DNA sequencing service with universal primers. Polymerase chain reaction (PCR) was performed with initial denaturation at 95°C for 2 min followed by 30 cycles with denaturation for 30 s at 94°C, annealing for 30 s at 58°C, and extension

for 45 s at 72 °C. The final extension was held for 5 min at 72 °C. The online program BLAST (NCBI, 2012) was used in identifying the related sequences with known taxonomic information available at the databank of National Center for Biotechnology Information (NCBI, Bethesda, Maryland, USA). A phylogenetic tree was constructed with the CLUSTAL X program¹⁷, which involved sequence alignment by the neighbor-joining method¹⁸ and maximum parsimony with the MEGA4 program¹⁹. The grouping of sequences was based on confidence values obtained by bootstrap analysis of 1000 replicates. Gaps were edited in the BioEdit program²⁰ and evolutionary distances were calculated by Kimura's two-parameter model²¹.

Qualitative estimation of phosphate solubilization

Phosphate solubilizing efficiency was assessed on PVK and NBRIP solid media^{22,23,24}. The isolate was spotted on solid media using sterile tooth picks, and plates were incubated at 30 ± 2°C for 2 days and the diameter of each colony as well as the halo zone formed were measured.

Quantitative estimation of soluble phosphate

For quantitative estimation of soluble P²⁴, a pre-culture of the bacterial isolate was prepared in 20 ml LB medium dispensed in 100 ml Erlenmeyer flask and incubated under shacked condition (170 rpm) at 30 ± 2°C till $\text{OD}_{600} = 1.0$ ($\sim 6 \times 10^9$ CFU/ml). For phosphate solubilization experiment, aliquots of 20 ml of either NBRIP or PVK were dispensed in 100 ml Erlenmeyer flasks and inoculated with 0.2 ml (1% v/v) of previously prepared seed cultures. Liquid cultures were incubated in a shaker incubator (170 rpm) for 5 days at 30 ± 2°C, unless otherwise stated. Uninoculated flasks were kept as control. The content of each flask was then centrifuged at 10,000 rpm for 10 min to remove cells and the supernatants were analyzed for soluble P content using the molybdenum-blue method²⁵. All experiments were carried in triplicates and the values are the average of data obtained.

Experimental Design and Optimization

Plackett-Burman design a rapid screening multifactor to find the most significant independent factors^{26,27,28}, was used in the present study to screen the important variables that significantly influenced phosphate solubilization. In this experiment, seven independent variables were screened in eight combination organized according

to the Plackett-Burman design matrix (Table2). Each variable was evaluated at two levels, a high (+) and a low (-) level (Tables 3 and 4.). All trials were performed in triplicates and the averages of phosphate solubilization (PS%) were treated as a response. Higher PS % is directly correlated with higher solubilization efficiency. This efficiency was evaluated based on PS%, all treatments were obtained at 12 h of inoculation.

The main effect of each variable can be calculated using the following standard equation:

$$\text{Main effect} = [\bar{R}(H) - \bar{R}(L)]/N$$

Where $R(H)$ and $R(L)$ are observations of trials where the independent variable was present in high and low concentrations, respectively, and N is the number of the trials divided by 2. A main effect figure with a positive sign indicates that the high concentration of this variable is near to optimum and a negative sign indicates that the low concentration of this variable is near to optimum.

Data Analysis of the Result of Plackett-Burman Design

Excel (Microsoft Office, 2010) was used for the experiment design and all statistical analyses. The variables with confidence levels above 95% were considered as influencing phosphate solubilization significantly. The application of statistical design was carried out in a 'two phases' of optimization approach. The first step was to evaluate the relative importance of various constituents within a complex culture medium and selecting of levels of the variables, which have significant influences on the solubilization process. The second was the verification experiments to validate the results under specific optimized experimental conditions.

RESULTS AND DISCUSSION

Identification of isolate H3fr

A total of 45 isolates that exhibited clear zones on the agar plates were selected as phosphate solubilizing organisms. One isolate (H3fr) was selected as an efficient phosphate solubilizing organism. The selected isolate had a marked insoluble phosphate solubilizing ability as visualized by the clear zone development around the colonies after 2 days of incubation. Comparison of the 16S rRNA sequence among available strains

of *Erwinia* species showed the highest sequence homology to *Erwinia* sp. CMG3059 EU048331.1 with the similarity percentage of 98.5% and to *Erwinia* sp. v427 EF088378.1 and *Enterobacter* sp. EU567059.1 with the similarity percentage of 98%. This strain was designated as *Erwinia* sp. h3fr. GenBank accession number of 16S rDNA gene partial sequence of the selected isolate and the highest sequence similarity as well as the closest neighbour(s) are shown (Table 1). Reference sequences were retrieved from GenBank under the accession numbers indicated in the trees. The obtained sequence was deposited in the NCBI Genebank under accession number GU182309. **Compared with all sequences, the results are presented as a phylogenetic tree (Fig. 1), showing the evolutionary position of H3fr amongst related organisms**

The genus *Erwinia* belongs to family *Enterobacteriaceae*, *Gamma Proteobacteria*. Indeed, several Gram-negative, non-spore-forming, facultatively anaerobic, rod-shaped bacteria have been classified traditionally into the genus *Erwinia*, mainly on the basis of their association with plants as pathogens, epiphytes or saprophytes^{29,30}.

In accordance with our data, it was reported that the majority of PSB isolates was affiliated to the *Beta* or *Gamma* sub-divisions of the *Proteobacteria*; three were similar to *Pantoea* species and two were closely related to *Serratia* species³¹. *Erwinia* was listed as one of the commonly reported inorganic phosphate mineralization genera^{32, 33, 34, 35, 36}. From a total of 395 rhizobacterial isolates, two *Erwinia* sp. (AUEY28 and AUEY29) were described as the top mineral P mobilizers isolated³⁶.

Qualitative and quantitative estimation of P solubilization efficiency by *Erwinia* sp. h3fr

Selection of suitable medium is an important issue. Data in Fig. 2 depict that medium NBRIP appeared to support the solubilization process for *Erwinia* sp. h3fr as observed by the clear zone formed. For quantitative measurement, standard inoculum (1%) of *Erwinia* sp. h3fr was used to inoculate 20 ml portion of NBRIP and PVK liquid media. Flasks were then incubated at 30°C for 5 days. As shown in Fig. 3 the value of soluble phosphorus (84.5%, 839 µg/ml) obtained with NBRIP was almost double the amount produced on PVK medium (42.2%, 418 µg/

ml). These data are in good agreement with those previously reported who found a maximum production of 450 µg/ml on NBRIP broth by isolate NBRIP12601^{37,38}. Similarly, the isolate EnHy-401 had the maximum value of soluble phosphorus (632.6 ± 23.0 µg/ml)³⁹ after 5 days incubation⁴⁰. Also, isolate BIHB752 recorded a maximum solubilization which liberated 805.0 µg/ml phosphorus⁴⁰. The values recorded in this study were higher than those

isolates from Korean soils that reported TCP solubilisation (96 to 139 (µg/ml)⁴¹. *Pantoea agglomerans* isolated from the acidic soils of Korea recorded 900 µg/ml soluble phosphorus⁴².

Under our experimental condition, *Erwinia* sp. hdds3fr was able to grow on the two tested media and solubilize tricalcium phosphate. As clearly observed the NBRIP medium supported better solubilization compared to PVK⁴¹. Therefore,

Table 1. Accession number of isolate H3fr and its similarity percentage to the nearest neighbours

Isolate	Genbank accession number	Similarity (%)	Nearest neighbour(s)
H3fr	GU182309	98.5	<i>Erwinia</i> sp. CMG3059 EU048331.1
		98	<i>Erwinia</i> sp. v427 EF088378.1
		98	<i>Enterobacter</i> sp. EU567059.1




Fig. 1. Phylogenetic tree of isolate H3fr. 16S rDNA-based dendrogram showing the phylogenetic position of isolate H3fr among representatives of related bacterial species. The tree was constructed by Bioedit method²⁰. Percentages (right) indicate phylogenetic similarities

further experimentation was done using NBRIP medium

Optimization of P solubilization by *Erwinia* sp.hdds3fr

For screening of the most important factors affecting PS, the Plackett-Burman design was applied with nine different fermentation conditions as shown in Table 3. All experiments were performed in triplicate and the average of results of phosphate solubilization (%) was




Fig. 2. Clear zone formation by *Erwinia* sp.hdds3fr grown for 48 h on PVK (A), and NBRIP (B) media

presented as response as shown in Table 3. The main effect of each variable on phosphate solubilization as well as t-values were estimated for each independent variable as shown in Table 4 and graphically presented in Figure 4. The values of PS% for different treatments showed a variation from 23 to 72% (Table 3), suggesting that the studied

variables had a significant effect on the PS% and therefore on phosphate solubilization.

Every microorganism has its own peculiar nutritional and physico-chemical requirements for the enzyme production⁴³; where the main effect with a positive sign indicates that the high concentration of this variable is nearly optimum




Fig. 3. Soluble P ($\mu\text{g ml}^{-1}$) in culture filtrate of *Erwinia* sp. hdds3fr as affected by different cultivation medium. Cultures were incubated at 30°C, 170 rpm for 5 days

Table 2. Factors examined as independent variables affecting phosphate solubilization by *Erwinia* sp. hdds3fr and their levels in the Plackett-Burman experimental design

Variable	Symbol	Level (g/l)		
		Lower	0 level	Higher
Glucose	G	5	10	15
$\text{Ca}_3(\text{PO}_4)_2$	C	2.5	5	7.5
MgCl_2	Mc	2.5	5	7.5
$\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$	Ms	0.125	0.25	0.375
KCl	K	0.1	0.2	0.3
$(\text{NH}_4)_2\text{SO}_4$	N	0.05	0.1	0.15
culture volume	V	10 ml	20 ml	30 ml

Table 3. The Plackett-Burman experimental design (in coded levels) with PS % as response

Trials	Factor under study	Component						Response [Soluble P (%)]	
		G	C	Mc	Ms	K	N		
1	Ms K N	- 1	- 1	- 1	+1	+1	+1	- 1	55
2	G N V	+1	- 1	- 1	- 1	- 1	+1	+1	72
3	C V	- 1	+1	- 1	- 1	- 1	- 1	+1	24
4	G C K	+1	+1	- 1	- 1	+1	- 1	- 1	23
5	McMs V	- 1	- 1	+1	+1	- 1	- 1	+1	50
6	G McMs K	+1	- 1	+1	+1	+1	- 1	- 1	52
7	C Mc N	- 1	+1	+1	- 1	- 1	+1	- 1	23
8	G C McMs K N V	+1	+1	+1	+1	+1	+1	+1	25
9	Basal medium	0	0	0	0	0	0	0	46






Fig. 4. Elucidation of culture conditions affecting phosphate solubilization (PS%) by *Erwinia* sp. hdds3fr

and a negative sign indicates that the low concentration of this variable is nearly optimum. In this study with respect to *Erwinia* sp. hdds3fr, the analysis of the main effect, t-values and P-values of 7 factors presented graphically (Fig. 4) show that the four variables G, MS, N and V positively affected phosphate solubilisation. Meanwhile, C, Mc and K had negative effects. Positive effect explains that if a higher concentration was used, a better response was

Table 4. Statistical analysis of the Plackett-Burman experimental result for phosphate solubilization by *Erwinia* sp. hdds3fr

Factor	Phosphate solubilization			
	Main effect	t-value	p-value	Significance level
Glucose	5.106	0.351461	0.74	26%
$\text{Ca}_3(\text{PO}_4)_2$	-33.779	-6.726512	0.01	99%
MgCl_2	-5.779	-0.398964	0.71	29%
MgSO_4	10.442	0.743560	0.49	51%
KCl	-3.271	-0.223790	0.84	16%
$(\text{NH}_4)_2\text{SO}_4$	6.716	0.465845	0.66	34%
Culture Volume	4.532	0.311251	0.77	23%

**Fig. 5.** Pareto plot of standardized effects**Fig. 6.** Verification experiment of the applied Plackett-Burman statistical design by comparing the percentage of soluble phosphorus in *Erwinia* sp.hdds3fr culture filtrate, growing on the optimized, basal and anti-optimized media

achieved, while a negative effect means lower concentrations are favored for better results.

Phosphate solubilisation was increased with the increase of glucose in the culture of *Erwinia* sp. hdds3fr as it was reported with *Burkholderia cepacia* DA23⁴⁴, *Acinetobacter calcoaceticus* YC-5a⁴⁵, *Serratia marcescens* CTM 50650⁴⁵, *Pantoea agglomerans* R-42⁴² and *Pseudomonas* sp.³⁸. The positive effect of magnesium sulphate in enhancing P solubilization observed in this study was also reported with *Acinetobacter calcoaceticus* YC-5a⁴⁴. It was reported that, the rate of phosphate solubilization by *Pseudomonas* sp. was increased with increasing concentrations of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ and MgCl_2 ; suggesting that, KCl is the less required ingredient and the phosphate solubilization ability improved when $(\text{NH}_4)_2\text{SO}_4$ was used at a lower concentration³⁸. On the contrary, in the present

study, P solubilisation required high concentration of $(\text{NH}_4)_2\text{SO}_4$. The presence of the insoluble tricalcium in low level enhanced the solubilization process. On the other hand and in contrast with *Erwinia* sp.hdds3fr; it was found that, the soluble P production was enhanced with increasing amount of insoluble phosphates by *Burkholderia cepacia* DA23 and *Acinetobacter calcoaceticus* YC-5a, respectively^{44,45}. From the data obtained, it was concluded that increasing the volume of culture medium (low aeration) might increase the amount of soluble phosphorus. In another study, it was found that phosphate solubilization by *Pantoea agglomerans* (KCTC 2564) increased independently of aeration⁴⁷, while phosphate removal by *Burkholderia* spp. from sediments containing mineral-phosphates was facilitated in the absence of aeration⁴⁸. On the contrary, in a study on phosphate mobilization by *Burkholderia*

caribensis FeGL03, it was found that restricted air supply in cultures would then adversely affect gluconic acid production⁴⁹.

The significant variables were identified by statistical analysis of the Plackett-Burman experiment using the t-test supported by Excel (Microsoft Office, 2010) to determine the statistical significance of the measured response (Table 4). Some researchers employed confidence levels greater than 70 % as significant effect levels^{50, 51}. Other investigators find that, confidence levels greater than 90 % are acceptable^{52, 53, 54} or even levels greater than 95 % to be a significant effect on the response^{55, 45}. In the present study, any component showing a statistical confidence equal and/or higher than 95 % ($P<0.05$) was considered significant parameter. It was clear that only one variable $[\text{Ca}_3(\text{PO}_4)_2]$ had a significant level ($P = 0.01$) with main effects of 33.779 %, while the other six variables with $P>0.05$, were considered insignificant.

Fig.5. shows the Pareto's chart that represents the estimated effects, in decreasing sequence of magnitude, of C, MS, N, Mc, G, V, and K. The measure of each bar is proportional to the estimate effect. Pareto's chart is a graphical representation of Student's t-test. This test presents each of the estimated effects. To plot the estimates in decreasing order of importance Pareto's chart is selected^{56, 52}.

Verification experiment

In order to validate the obtained data and to evaluate the accuracy of the applied Plackett-Burman statistical design, a verification experiment was carried out in triplicates. Consequently, based on the results obtained from the Plackett-Burman experiment, a medium of the following formula was predicted to be near optimum for phosphate solubilization (g/l): glucose, 15; $\text{Ca}_3(\text{PO}_4)_2$, 2.5; MgCl_2 , 2.5; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.375; KCl , 0.1; $(\text{NH}_4)_2\text{SO}_4$, 0.15 and culture volume, 30 ml. The data were examined and compared to the basal and anti-optimized medium. The anti-optimized medium was the opposite concentrations of the optimized one and its formula was in (g/l): glucose, 5; $\text{Ca}_3(\text{PO}_4)_2$, 7.5 ; MgCl_2 , 7.5; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.125; KCl , 0.3; $(\text{NH}_4)_2\text{SO}_4$, 0.05 and culture volume, 10 ml.

The verification experiment applied indicated that the solubilized P before and after optimization were 46% and 90%, respectively. Thus,

a significant increase near 2 fold in SP was achieved by Plackett-Burman optimization (Fig. 6). On the other hand, *Acinetobacter calcoaceticus* YC-5ashowed 1.8 fold phosphate solubilisation increase compared with that using the original medium⁴⁵.

CONCLUSION

The results presented here emphasize the importance of exploring the natural biodiversity of local environments and studying P-solubilizing microorganisms. Data also demonstrate optimum metabolic capacity at different environmental parameters by the application of statistical experimental design that would probably enhance the solubilization process. It is suggested that researchers must continue to examine microbial interactions for use as biofertilizers, and growth promoters in order to improve communities in field soils.

REFERENCES

1. Khan, M. S., Zaidi, A., Ahmad, M., Oves, M., Wani, P.A. Plant growth promotion by phosphate solubilizing fungi – current perspective. *Arch. Agron. Soil Sci.* 2010; **56**: 73–98.
2. Sharma, S. B., Sayyed, R. Z, Trivedi, M. H., Gobi, T. A. Phosphate solubilizing microbes: sustainable approach for managing phosphorus deficiency in agricultural soils. *Springer Plus.* 2013; **2**: 587.
3. Saber, K., Nahla, L.D., Chedly, A. Effect of P on nodule formation and N fixation in bean. *Agron Sustain Dev.* 2005; **25**: 389–393.
4. Rengel, Z., Marschner, P. Nutrient availability and management in the rhizosphere: exploiting genotypic differences. *New Phyto.* 2005; **168**: 305–312.
5. Illmer, .P. A., Schinner, F. Solubilization of inorganic calcium phosphates solubilization mechanisms. *Soil Biol. Biochem.* 1995; **27**: 257–263.
6. Norrish, K., Rosser, H. Mineral phosphates. In *Soils: An Australian Viewpoint*. CSIRO Division of Soils, Academic Press. 1983; pp 335–364.
7. Lindsay, W. L., Vlek, P. L. G., Chien, S. H. Phosphate minerals. In: *Minerals in soil environment* (Dixon JB, Weed SB, eds), 2nd edn. Soil Science Society of America, Madison, WI,

USA.1989; pp 1089–1130.

8. Khan, A.A., Jilani, G., Akhtar, M.S., Naqvi, S.M.S., Rasheed, M. Phosphorus solubilizing bacteria: occurrence, mechanisms and their role in crop production. *J. Agric. Biol. Sci.* 2009a; **1**:48–58.
9. Khan, M.S., Zaidi, A., and Wani, P.A. Role of phosphate solubilizing microorganisms in sustainable agriculture. In: *Sustainable Agriculture* (Lictfouse E et al., eds). Springer. 2009b; pp 552, DOI: 10.1007/978-90-481-2666-8_34.
10. Khan, M.S., Zaidi, A., Wani, P.A., Oves, M. Role of plant growth promoting rhizobacteria in the remediation of metal contaminated soils. *Environ. chem. let.* 2009c; **7**:1–19.
11. Cordell, D., Drangert, J.O., White, S. The story of phosphorus: global food security and food for thought. *Glob. Environ. Chang.* 2009; **19**: 292–305.
12. Isherwood, K.F. Mineral Fertilizer Use and the Environment. International Fertilizer Industry Association/United Nations Environment Programme, Paris. 2000.
13. Zaidi, A., Khan, M.S., Ahmad, M., Oves, M., Wani, P.A. Recent Advances in Plant Growth Promotion by Phosphate-Solubilizing Microbes. In: *Microbial Strategies for Crop Improvement* (Khan, M.S., Zaidi, A., Musarrat, J., eds). Springer-Verlag, Berlin Heidelberg. 2009; pp 23–50.
14. Johri, J.K., Surange, S., Nautiyal, C. S. Occurrence of Salt, pH, and Temperature-tolerant, Phosphate-solubilizing Bacteria in Alkaline Soils. *Curr.Microbiol.* 1999; **39**: 89–93.
15. Turan, M., Ataoglu, N., Sahin, F. Evaluation of the capacity of phosphate solubilizing bacteria and fungi on different forms of phosphorus in liquid culture. *J Sustain. Agr.* 2006; **28**: 99–108.
16. Vyas, P., Rahi, P., Gulati, A. Stress tolerance and genetic variability of phosphate-solubilizing fluorescent *Pseudomonas* from the cold deserts of the trans-Himalayas. *Microb. Eco.* 2009; **58**:425-434.
17. Thompson, J.D., Gibson, T.J., Plewniak, F., Jeanmougin, F., Higgins, D.G. The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* 1997; **24**: 4876–4882.
18. Saitou, N., Nei, M. The Neighbor-Joining method—a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* 1987; **4**: 406–425.
19. Kumar, V., Behl, R.K., Narula, N. Establishment of phosphate-solubilizing strains of *Azotobacter chroococcum* in the rhizosphere and their effect on wheat cultivars under greenhouse conditions. *Microbiol. Res.* 2001; **156**: 87–93.
20. Hall, T.A. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucl. Acids. Symp. Ser.* 1999; **41**: 95–98.
21. Kimura, M. A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *J. Mol. Evol.* 1980; **16**: 111–120.
22. Kundu, B. S., Kiran, N., Yadav, R., Tomar, M. Biodiversity of phosphate solubilizing bacteria in rhizosphere of chickpea, mustard and wheat grown in different regions of Haryana. *Ind. J. Microbiol.* 2009; **49**: 120–127.
23. Hu, X., Li, Z., Cao, Y., Zhang, J., Gong, Y., Yang, Y. Isolation and identification of a phosphate-solubilizing bacterium *Pantoeastewartii* subsp. *stewartii* g6, and effects of temperature, salinity, and pH on its growth under indoor culture conditions. *Aquacult. Int.* 2010; **18**: 1079–1091.
24. Ranjan, A., Mahalakshshmi, M.R., Sridevi, M. Isolation and characterization of phosphate-solubilizing bacterial species from different crop fields of Salem, Tamil Nadu, India. *Int. J. Nutri. Pharm. Neurol. Dis.* 2013; **3**: 29–33.
25. Murphy, J., Riley, J.P. A modified solution method for the determination of phosphate in natural waters. *Anal. Chim. Acta.* 1962; **27**: 31–36.
26. Plackett, R.L., Burman, J.P. The design of optimum multifactorial experiments. *Biomet.* 1946; **33**: 305–25.
27. Wang, X., Yin, M., Xiao, Z., Ma, C., Lin, Z., Wang, P.G., Xu, P. Production of uridine 5'-monophosphate by *Corynebacterium ammoniagenes* ATCC 6872 using a statistically improved biocatalytic process. *Appl. Microbiol. Biotechnol.* 2007; **76**: 321–328.
28. Xiao, Z., Liu, P., Qin, J., Xu, P. Statistical optimization of medium components for enhanced acetoin production from molasses and soybean meal hydrolysate. *Appl. Microbiol. Biotechnol.* 2007; **74**: 61–68.
29. Lelliott, R. A., Dickey, R. S. Genus VII. *Erwinia* Winslow, Broadhurst, Buchanan, Krumwiede, Rogers and Smith 1920. Baltimore: Williams & Wilkins. In *Bergey's Manual of Systematic Bacteriology* (Krieg, N. R. and Holt, J. G. eds.). 1984; **1**: 469–476.
30. Waleron, M., Waleron, K., Podhajska, A.J., Dojkowska E. Genotyping of bacteria belonging to the former *Erwinia* genus by PCR-RFLP analysis of a *recA* gene fragment.

31. *Microbiol.* 2002; **148**: 583–595.

32. Pe'rez, E., Sulbarán, M., Ball, M.M., Yarzábal, L.A. Isolation and characterization of mineral phosphate-solubilizing bacteria naturally colonizing a limonitic crust in the south-eastern Venezuelan region. *Soil Biol. Biochem.* 2007; **39**: 2905–2914.

33. Rodriguez, H., Fraga, R. Phosphate solubilizing bacteria and their role in plant growth promotion. *Biotech. Adv.* 1999; **17**: 319–339.

34. Ahmed, N., Shahab, S. Phosphate Solubilization: Their Mechanism Genetics And Application. *The Int. J. Microbiol.* 2011; **10**: 5580/2327.

35. Zhu, F., Qu, L., Hong, X., Sun, X. Isolation and characterization of a phosphate-solubilizing halophilic bacterium *Kushneria* sp. YCWA18 from DaqiaoSaltern on the coast of Yellow Sea of China. *Evid Based Complement Alternat Med.* 2011; **10**: 1155/615032.

36. Hamdali, H., Moursalou, K., Tchangbedji, G., Ouhdouch, Y., Mohamed, H. Isolation and characterization of rock phosphate solubilizing actinobacteria from a Togolese phosphate mine. *Afr. J. Biotechnol.* 2012; **11**: 312-320.

37. Muleta, D., Assefa, F., Börjesson, E., Granhall, U. Phosphate- Solubilising rhizobacteria associated with *Coffea arabica* L. in natural coffee forests of south-western Ethiopia. *J. Saudi. Soc. Agri. Sci.* 2013; **12**: 73–84.

38. Nautiyal, C.S., Bhaduria, S., Kumar, P., Lal, H., Mondal, R., Verma, D. Stress induced phosphate solubilization in bacteria isolated from alkaline soils. *FEMS Microbiol. Lett.* 2000; **182**: 291–296.

39. Nautiyal, C. An efficient microbiological growth medium for screening phosphate solubilizing microorganisms. *FEMS Microbiol. Lett.* 1999; **170**: 265–270.

40. Yi, Y., Huang, W., Ge, Y. Exopolysaccharide: a novel important factor in the microbial dissolution of tricalcium phosphate. *World J. Microb. Biotechnol.* 2008; **24**: 1059–1065.

41. Gulati, A., Rahi, P., Vyas, P. Characterization of phosphate- solubilizing fluorescent pseudomonads from the rhizosphere of seabuckthorn growing in the cold deserts of himalayas. *Curr. Microbiol.* 2008; **56**: 73–79.

42. Chung, H., Park, M., Madhaiyan, M., Seshadri, S., Song, J., Cho, H., Sa, T. Isolation and characterization of phosphate solubilization bacteria from the rhizosphere of crop plants of Korea. *Soil Biol. Biochem.* 2005; **37**: 1970–1974.

43. rhizosphere. *Biores. Technol.* 2006; **97**: 204–210.

44. Kumar, A.G., Nagesh, N., Prabhakar, T.G., Sekaran, G. Purification of extracellular acid protease and analysis of fermentation metabolites by *Synergistes* sp. utilizing proteinaceous solid waste from tanneries. *Biores. Technol.* 2008; **99**: 2364–2372.

45. Song, O., Lee, S., Lee, Y., Lee, S., Kim, K., Choi, Y. Solubilization of insoluble inorganic phosphate by *Burkholderiacepacia*DA2 isolated from cultivated soil. *Brazil J. Microbiol.* 2008; **39**: 151–156.

46. Fan, D. D., Ren, Y. X., Zhu, X. L., Ma, P., Liang, L. H. Optimization of culture conditions for phosphate solubilizing by *Acinetobacter calcoaceticus*YC-5a using response surface methodology. *Afr. J. Microbiol. Res.* 2011; **5**: 3327–3333.

47. Farhat, M.B., Farhat, A., Bejar, W., Kammoun, R., Bouchaala, K., Fourati, A., Antoun, H., Bejar, S., Chouayekh, H. Characterization of the mineral phosphate solubilizing activity of *Serratia marcescens* CTM 50650 isolated from the phosphate mine of Gafsa. *Arch. Microbiol.* 2009; **191**: 815–824.

48. Jung, P. D., Park, K. A Study of the growth condition and solubilization of phosphate from Hydroxyapatite by *Pantoea agglomerans*. *Biotechnol. Bioprocess Eng.* 2002; **7**: 201–205.

49. Kim, Y.H., Bae, B., Choung, Y.K. Optimization of biological phosphorus removal from contaminated sediments with phosphate solubilizing microorganisms. *J. Biosci. Bioeng.* 2005; **99**: 23–29.

50. Delvasto, P., Ballester, A., Muñoz, J.A., González, F., Blazquez, M.L., Igual, J.M., Valverde, A., García-Balboa, C. Mobilization of phosphorus from iron ore by the bacterium *Burkholderiacaribensis* FeGL03. *Min. Eng.* 2009; **22**: 1–9.

51. Stowe, R.A., Mayer, R.P. Efficient screening of process variables. *Ind. Eng. Chem.* 1966; **58**: 36–40.

52. Reddy, M.B.V. R., Rao, C.S., Rao, S.S.L. Evaluation of Process Parameters and Media Components by Plackett-Burman Design for Enhancement of Biomass using Cyanobacteria (*Anabaena ambigua*). *Int. J. Chem. Tech. Res.* 2012; **4**: 761–766.

53. Chen, Q.H., He, G.Q., Ali, M.A.M. Optimization of medium composition for the production of elastase by *Bacillus* sp. EL31410 with response surface methodology. *Enz. Microb. Technol.* 2002; **30**: 667–672.

54. AbdelFattah, Y.R., Olama, Z.A. L-Asparaginase production by *Pseudomonas aeruginosa* in

solid-state culture: evaluation and optimization of culture conditions using factorial designs. *Pro.Biochem.* 2002; **38**: 115-122.

54. Singh, A.K., Mehta, G., Chhatpar, H.S. Optimization of medium constituents for improved chitinase production by *Paenibacillus* sp. D1 using statistical approach. *Lett. Appl. Microbiol.* 2009; **49** :708–714.

55. Niladevi, K.N., Sukumaran, R.K., Jacob, N., Anisha, G.S., Prema, P. Optimization of laccase production from a novel strain-*Streptomyces psammoticus* using response surface methodology. *Microbiol.Res.* 2009; **164**: 105-113.

56. Abdel Fattah, Y.R. Optimization of thermostable lipase production from a thermophilic *Geobacillus* sp. using Box-Behnken experimental design. *Biotechnol. Lett.* 2002; **24**: 1217-1222.

© The Author(s) 2016. **Open Access**. This article is distributed under the terms of the [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) which permits unrestricted use, sharing, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.