

A First DNA Sequencing of Hydatid Agent Isolated from Human in Iraq

Abdulsadah A. Rahi^{1*} and Magda A. Ali²

¹Department of Biology, College of Science, Wasit University, Kut, Iraq.

²Department of Microbiology, College of Medicine, Wasit University, Kut, Iraq.

<https://doi.org/10.22207/JPAM.10.2.19>

(Received: 25 March 2016; accepted: 09 May 2016)

A total of 20 human hydatid cysts fluid samples were collected during January, 2015 to July, 2015 from patients attended to Al- Karamah teaching hospital at Wasit Province. Genetic characterization of all isolates of *Echinococcus granulosus* from hydatid cyst was performed by sequencing analysis of fragment of mitochondrial (CO1) cytochrome C oxidase subunit 1 and NADH dehydrogenase subunit1 (ND1) genes to determine genotyping of *E.granulosus* in hyper-endemic areas of Wasit province. This work done by randomly choosing of five samples. The results of DNA sequencing showed genotype 1(G1)and genotype 3 (G3) and the sheep strain G1was more prevalent than buffalo strain G3.

Keywords: Hydatid , Human, DNA sequence, Diagnosis.

Echinococcosis, or Hydatid disease, is an infection caused by tapeworms of the genus *Echinococcus*, a tiny tapeworm just a few millimetres long. Echinococcosis is a zoonosis, a disease of animals that affects humans. This disease is still a major endemic, zoonotic and public health problem also is not yet an organized national control program in Iraq¹. Outbreak of this disease have occurred in Asian countries such as Lebanon, Jordan, Iraq, Saudi Arabia and Iran leading to substantial health problems and economic losses². This parasite have five medical important species of genus *Echinococcus* are *E.granulosus*, *E. multilocularis* , *E. vogeli* , *E. oligarthrus* and *E. shiquicus* , these species are morphologically distinction both adult and larval stages³.

To date ten distinct genotype (G1-G10) have been described in world based on nucleotide sequence analysis of mitochondrial cytochrome oxidase subunit 1(CO1) (4). It has been reported that at least seven of these strains were isolated

from human, but the most common one is sheep strain^{4,5}.

The cysts of *Echinococcus granulosus* found in the most people one cyst , approximately(60-70%)of *Echinococcus granulosus* cysts in the liver and (20-25%) in the lungs⁶, also found these cysts in the other organs of human including “the bones, kidneys, spleen, muscles , central nerve system (CNS) and behind the eyes⁷.

In human echinococcosis is diagnosed mainly with imaging techniques such as ultrasonic graph, radiology, magnetic resonance imaging (MRI) or (T-scanning) supported by serology tests⁸. Serological tests used in humans include enzyme linked immune-sorbent assays (ELISAs), indirect immune fluorescence, indirect hemagglutination, immune blotting and latex agglutination^{9,10}. Molecular diagnosis such as DNA sequencing technique to determine the genetic characterization of *E.granulosus* depends on the homology of the sequence of two mitochondrial genes: cytochrome C oxidase subunit 1(CO1) and reduced nicotinamide adenine dinucleotide subunit 1(ND1)^{4,11}.

* To whom all correspondence should be addressed.
E-mail: abdulsadah1966@yahoo.com

MATERIALS AND METHODS

Samples collection

A total of 20 human hydatid cyst fluid samples were collected during January, 2015 to July, 2015 from patients attended to Al- Karamah teaching hospital at Wasit province and placed in sterile container, then transported to laboratory and stored in freeze (-20C) until genomic DNA extraction step.

Direct examination

The fluid of hydatid cyst was taken from patient after surgical immediately and centrifuged. The precipitate smeared on clean slide and stained by Eosin (0.1%), then examined by microscope under (40x) to show the protoscolices clearly.

DNA Extraction

Genomic DNA from hydatid cyst fluid samples was extracted by using Genomic DNA mini kit extraction Geneaid, USA, and performed according to company's instructions.

DNA Profile

The extracted DNA was checked by Nanodrop spectrophotometer (THERMO, USA), which measured DNA concentration (ng/ μ L) by reading the absorbance at (260 /280 nm).

PCR technique

PCR technique was performed for detection of two mitochondrial genes in *E. granulosus* that using in genotyping study from human samples. This technique was carried out according to method described by¹² and provided by Bioneer company, Korea .

DNA sequencing method

DNA sequencing method was performed for genotyping and phylogenetic analysis study of local *Echinococcus granulosus* isolates based mitochondrial cox1 and nad1 gene. The sequencing

of the PCR product 450 bp and 400 bp for cox1 and nad1 gene respectively, where the PCR product was purified from agarose gel by using (EZ-10 Spin Column DNA Gel Extraction Kit, Biobasic, Canada). Phylogenetic analysis was performed based on NCBI-Blast Alignment identification and Unweighted Pair Group method with Arithmetic Mean (UPGMA tree) in (MEGA 6.0 version).

RESULTS AND DISCUSSION

Amplification of genes by PCR

Amplification of mitochondrial CO1gene

DNA was amplified by two primers for detection and genotyping of *E.granulosus* based on sequencing of mitochondrial genes by using PCR technique. These primers generated the expected (450 bp) product. The results showed all samples had CO1 gene (Figure 1).

Amplification of mitochondrial ND1gene

DNA was showed (NAD1) dehydrogenase subunit 1 gene amplified from all samples of human hydatid cysts by using specific primers . PCR product were purified after a garose gel electrophoresis at (400) bp and visualized under UV transilluminator (Figure 2).

Genotyping of *E.granulosus* isolates

In this study, genetic characterization of 20 isolates of *E.granulosus* from hydatid cyst was performed by sequencing analysis of fragment of mitochondrial CO1 gene and ND1 gene to determine genotyping of *E.granulosus* in hyper-endemic areas of Wasit province. This work done by randomly choosing of five samples. The results of DNA sequencing showed genotype 1 (G1) and genotype 3 (G3) (Figure 3,4). This is the first study at Wasit province, Iraq (Figures 3,4,5).



Fig. 1. Agarose gel electrophoresis image shown the PCR product analysis of cox1 gene in *E.granulosus* , Where M marker (2000-100bp), lane (1-10) positive cox1 gene at 450 bp PCR product.




Fig. 2. Agarose gel electrophoresis image shown the PCR product analysis of nad1 gene in *E.granulosus* samples, Where M marker (1500-100bp), lane (1-10) positive nad1 gene at 400 bp PCR product




Fig. 3. Phylogenetic tree analysis based on the cox1 gene of *E.granulosus*. The isolates (1, 3, 4, 5) were shown closely related to NCBI-Blast *E. granulosus* isolate (Genotype 1) (KF612390.1), whereas the isolate (2) was shown closely related to NCBI-Blast *E. granulosus* isolate (Genotype 3) (KF612397.1)




Fig. 4. Phylogenetic tree analysis based on the nad1 gene of *E.granulosus*. The isolates (1,3,4,5) were shown closely related to NCBI-Blast *E. granulosus* isolate (Genotype 1) (KF612360.1), whereas the isolate (2) was shown closely related to NCBI-Blast *E. granulosus* isolate (Genotype 3) (KF612369.1)




Fig. 5. Multiple sequence alignment analysis of the partial cox1 gene sequence for local *E. granulosus* isolates and NCBI-Genbank *E. granulosus* isolates. The multiple alignment analysis similarity (*) and differences in cox1 gene nucleotide sequences




Fig. 6. Multiple sequence alignment analysis of the partial nad1 gene sequence for local *E. granulosus* isolates and NCBI-Genbank *E. granulosus* isolates. The multiple alignment analysis similarity (*) and differences in nad1 gene nucleotide sequences




Fig. 7. A,B. Phylogenetic tree analysis based on the Cox1 gene (A) and nad1 gene (B) partial sequence used for *E. granulosus* country relationship. The local *E. granulosus* isolates (S1, S3 and S4) shown closely related to neighboring countries available NCBI-Blast *E. granulosus* genotype 1 Iran isolates (JQ250810.1) and (JQ219962.1), Whereas other local *E. granulosus* isolate (S5) shown closely related to NCBI-Blast *E. granulosus* genotype 1 China isolate (AB688615.1)

DISCUSSION

This is the first study to determine the genetic characterization of the strains of *E. granulosus* from humans at Wasit province. Two

genotypes G1 and G3 were identified among our DNA samples based on (CO1) and (ND1) sequences data, genotype G1 was the most common available genotype in the studied population. The phylogenetic analysis of concatenated sequences

of COX1 and NAD1 showed 4 samples (s1,s3,s4,s5) represent all strains related to G1 genotype and sample(s2) represents strain of G3 genotype.

The preliminary studies by¹³, the G1 was found in all humans isolated, also confirmed by study of¹⁴ , in Ilam province,Iran which reported that all of humans isolates belonged to G1 genotype. The disagreement of these results could be explained by type of studied samples, difference in geographic area and exploited method for genotyping¹⁵.

Numerous studies have been carried out on genetic characterization of hydatid cyst,using hydatid fluid, cystic membrane and scolices¹⁶.Various PCR techniques have been introduced to detect *E. granulosus* genome in biological specimens¹⁷. PCR is currently developed as acomplementary diagnostic tool for species differentiation for echinococcosis,using biopsy samples and tissue specimens^{18,19}.

The DNA sequencing of specific region of NADH Nicotinamide adenine dinucleotide subunit1gene confirmed the presence of sheep strain G1 and buffalo strain G3 at Wasit Province. When comparing the results of sequencing of ND1 in G1 and G3 by Gen-Bank were showed identity (100%) in Wasit province and identically with Iraqi also have identity (100%) for ND1 of G1 and G3. These result were comparatively closer to similar studies conducted by many authors in Iran²⁰, France²¹ and Romania²². However, different results were obtained by other authors worldwide²³.

Abdullah *et al.*, 2012²⁴ showed that sheep strain G1 is predominant in Kurdistan-Iraq and it was mostly responsible of human hydatid disease and Kia *et al.*, 2010²⁵, confirmed that G1 was the predominant genotype of hydatidosis in human in central Iran. As well as Eryildis and Sakra, 2012²⁶, proven that G1 was the predominant in Turkey and is the principle agent of human and animals hydatidosis. These agreed with our results that confirmed G1 predominant than G3 at Wasit Province .

The previous studies in the world were used mitochondrial and nuclear DNA markers in molecular characterization of *E.granulosus*, but have no adequate studies to determine and identify *E.granulosus* strains that are cause infection of human²⁷⁻²⁹. The recent studies reported that at

least seven of these strains are infective to human but the sheep strain and the G1 were the most common of these strains³⁰.

REFERENCES

1. Athmar, K.A.A and Ban-Abbas, A.M. Immunization mice with DNA from protoscolices of human hydatid cyst. A immunological study. *Int J Adv Bio Res.* 2014 ; **4**(1):89-95.
2. Abdi, J. kazemi, B. Karimfar, M. and Rokni, M. Evaluation of rabbit of antibody response against 8 and 16 KDa recombinant sub units of antigen B from *Echinococcus granulosus*. *Asian pac. J. Trop. Med.* 2012; 355- 357.
3. Abbasi, I. Branzburg Campos-Ponce, M. Abdel hafezs, K. Raoul, F. Craig, P.S. and Hamburger , J. Corpo-diagnosis of *Echinococcus granulosus* infection in dogs by amplification of a newly identified repeated DNA sequence .*Am.J.Trop.Med.Hyg.*2003; **69**: 324-330.
4. SaHchez, E. Caceres, O. Naquira, C. Garcia, D. Patino, G. Siliva, H. Volotao, A. C and Fernandes, O. Molecular characterization of *Echinococcus granulosus* from peru by sequencing of the mitochondrial cytochrome C Oxidase subunit 1 gene. 2010; **105** (6):806-810.
5. Eckret, J. Conraths, F.J and Tackmann, K. *Echinococcosis* : An emerging or R- emerging zoonosis.*Int .J. Parasitol.* 2000; **30**(12-13): 1283- 1294.
6. Foreyt, W.J. Drew, M. L. Atkinson, M. and Mccauley, D. *E.granulosus* in gray wolves and ungulates in Idaho and Montana, *USA. J wild. L. Dis.*2009; **45**(4):1208-1212.
7. Goto,Y. Satok Y. ahagi, K. Komatsu,O. Hoshina, H. Abiko, C. Yamasaki, H. and Kawanaka, M. Frequent isolation of *E. multilocularis* from the liver of races horses slaughtered in Yamagata , Japan. *Jpn .J. Infect Disease.* 2010; **63**(6):449- 451.
8. Dakkak , A. Echinococcosis / Hydatidosis : asereve threat in Mediterranean countries .*Vet parasitol.* 2010; **174**(1-2):2-11.
9. Roberts, L.S. and Janovy, J.J. Foundations of Parasitology. 6th ed. McGraw-Hill Companies, Inc. New York: 2000; 338-341.
10. Siracusano, A. Delunardo, F. Teggi, A. and Ortona, E. Host-parasite Relationship in Cystic Echinococcosis: An Evolving Story. *Hindawi Publishing Corporation, Clinical and Developmental Immunology*: 2012;1-12.
11. Snbel,V. Altintas, N. D,Amelio, S. Nakao, M. Romig, T. Yolasigmaz , A. Gunes, K. Turk, M.

Busi, M. Hqttner, M. Sevcov, D. Ito, A. Altintas, N. and Dubinsky, P. Cystic echinococcosis in Turkey: genetic variability and first record of the pig strain(G7) in the country. *parasitol.Res.* 2009; **105**(1): 145-154.

12. Nikmanesh, B. Hossein, M. Zohreh, G. Masoud, A. Mitra, S. Eshrat, B. K. Mehdi, M. Maryam, E. and Mohammad, B. Genotyping of *Echinococcus granulosus*. Isolates from Human Clinical Samples Based on Sequencing of Mitochondrial Genes in Iran. *Iran J Parasitol.* 2014; **9**(1):20-27.

13. Zhang, L. Eslami, A. Hosseini, S.H. and McManus, D.P. Indication of the presence of two distinct strains of *Echinococcus granulosus* in Iran by mitochondrial DNA markers. *Am J Trop Med Hyg.* 1998; **59**:171-175.

14. Dousti, M. Abdi, J. Bakhtiyari, S. Mohebali, M. Mirhendi, S.H. and Rokni, M. Genotyping of hydatid cyst isolated from human and domestic animals in Ilam province, western Iran using PCR-RFLP. *Iran J Parasitol.* 2013; **8**(1): 47-52.

15. Gholami, S.H. Sosari, M. Fakhar, M. Sharif, M. Daryani, A. Hashemi, M. Molecular Characterization of *Echinococcus granulosus* from Hydatid Cysts Isolated from Human and Animals in Golestan Province, North of Iran. *Iranian J. Parasitol.* 2012; **7**(4):8-16.

16. Dinkel, A. Njoroge, E. M. Zimmermann, A. Wälz, M. Zeyhle, E. Elmahdi, I. E. Mackenstedt, U. and Romig, T. A PCR System for Detection of Species and Genotypes of the *Echinococcus granulosus*-Complex, with Reference to the Epidemiological Situation in Eastern Africa, " *Int J Parasitol.* 2004; **34**(5): 645-653.

17. Siles-Lucas, M.M. and Gottstein, B.B. Molecular Tools for the Diagnosis of Cystic and Alveolar Echinococcosis. *Trop Med Int Hlth.* 2001; **6**(6): 463-475.

18. Kern, P. Frosch, P. Helbig, M. Wechsler, J. G. Usadel, S. Beckh, K. Kunz, R. Lucius, R. and Frosch, M. Diagnosis of *E. multilocularis* Infection by Reverse-Transcription Polymerase Chain Reaction. *Gastroenterology.* 1995; **109**(2): 596-600.

19. Myjak, P. Nahorski, W. Pietkiewicz, H. Von Nickisch-Rosenegk, M. Stolarczyk, J. Kacprzak, E. Felczak- Korzybska, I. Szostakowska, B. and Lucius, R. Molecular Confirmation of Human Alveolar Echinococcosis in Poland. *Clin Inf Dis.* 2003; **37**(8): 121-125.

20. Hajialio, E. Harandi, M.F. Sharbatkhori, M. Mirhendi, H. Rostami, S. Genetic characterization of *E.granulosus* in camels, cattle and sheep from the south east of Iran indicates the presence of the G3 genotype. *J*

Helminthes. 2011; **13**(86):1-8.

21. Umhang, G. Richomme, C. Boucher, J-M. Hormaz, V. Boué, F. Prevalence survey and first molecular characterization of *Echinococcus granulosus* in France. *Parasitol Res.* 2012; **112**: 1809-1812.

22. Stefania, S. Anamaria, I. Monica, J. and Vcozma. Prevalence of *E.granulosus*. Cluj county , Romania, revealed by PCR . *Science parasite J.* 2009; **1**(2):68-71.

23. Amir, V. Mahdi, M. Ardavan, G. and Behroz, S. Genotypic Characteristics of hydatid cysts isolated from humans in East Azerbaijan Province (2011-2013). *J. Anal. Res. Clin. Med.* 2013; **2**(3):152-157.

24. Abdullah, A. H. Wijda, M.S. M. and Jaladet, M.S. J. Molecular Characterization of *E.granulosus* First Report of Sheep Strain in Kurdistan-Iraq. 2nd international conference on Ecological, Enviromental and biological Science. 2012; 41-44.

25. Kia, E.B. Rahimi, H. Sharbitkhori, M. Talebi, A. Harandi, M.F. and Mirhendi, H. Genotype identification of human cystic echinococcosis in Isfahan, central Iran. *Parasitol. Res.* 2010 ; **107**:757-760.

26. Eryildis, C. and Sakru, N. Molecular characterization of human and animal isolates of *E.granulosus* in the thrace region, Turkey. *Balkan Med. J.* 2012; **29** (3): 261-267.

27. Karimi, A. and Dianatpour, R. Genotypic and phenotypic characterization of *Echinococcus granulosus*. *Boil.* 2008; **7**(4): 757-762.

28. Thompson, C.R.A. and Mcmanus, D.P., Aetiology parasites and life cycles . In: Eckert, J. Gemmell, M.A. Meslin, F. X. and Pawolowski, Z .S (Eds). WHO/OIE manual on echinococcosis in human and animals : A public health problem of global concern. *world organization for animal health.* 2001 Paris : 1-19.

29. Bhattacharya, D. Bera, A.K. Bera, B.C. Maity, A. and Das, S.K. Genotypic characterization of Indian cattle, Buffalo and sheep isolate of *E. granulosus* . *Vet. Parasitol.* 2007; **143**: 371-374.

30. Sharbatkhori, M. Mirhendi, H. Harandi, M. F. Rezaeian, M. Mohebali, M. Eshraghian, M. Rahimi, H. and Kia, E .B. *Echinococcus granulosus* genotypes in livestock of Iran indication high frequency of G1 genotype in Camels .*Exp. parasitol.* 2010; **124**(4): 373-379.

© The Author(s) 2016. **Open Access.** This article is distributed under the terms of the [Creative Commons Attribution 4.0 International License](http://creativecommons.org/licenses/by/4.0/) which permits unrestricted use, sharing, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.