

## Nanotechnology and Agriculture: A review

**Mukesh Srivastava, Sonika Pandey, Mohammad Shahid, Anuradha Singh, Vipul Kumar, Shubha Trivedi, Manoj Kumar Maurya, Yatindra Kumar Srivastava, Ved Ratan and D.K. Srivastava\***

Chandra Shekhar Azad University of Agriculture & Technology, Kanpur - 208 002, India.

\*Council of Science and Technology, Lucknow, India.

<https://doi.org/10.22207/JPAM.10.2.25>

(Received: 09 December 2015; accepted: 10 February 2016)

Nanotechnology is the most developing branch of science, it is thought that application of nanotechnology in agriculture will make this sector sustainable and profitable, by reducing the use of fertilizers, pesticides and by enhancing the crop productivity. In other words we can say that application of nanotechnology in agriculture will increase food and nutritional security. However, the main issue related with the application of nanoparticles in agriculture is their toxicity. It is therefore recommended for research organizations to provide research funds to establish safety standards and to develop new techniques for the toxic free synthesis of nanoparticles. Indian council of agriculture research (ICAR) has started a programme to study the effect of nanotechnology in agriculture. The ICAR- nanotechnology scheme has major themes such as synthesis of nano-particles for agriculture use, diagnostic kits for easy and quick identification of disease and pests, nano agri-inputs for enhanced use efficiencies, precision water management, and stabilization of organic matter in soil, nano food systems and bio safety besides establishing the policy frame work. Green synthesis and microbial synthesis of nanomaterial for agriculture use is very important as they are naturally encapsulated with mother protein, therefore more stable and safer to biological systems. The review presents here discuss applications of nanoparticles in agriculture. This review also features out the potential hazardous effects of nanoparticles on microbes and environment.

**Keywords:** Carbon nanotubes, Nanoparticles, Agriculture, Microbes.

Food crisis is a global problem, especially for developing countries. Even though India is agriculture based country but it also suffers from food crisis. There are reports which clearly show that India is facing a steady decline in its GDP. India had made a record of producing 241 million tones of crop production in 2010-2011([www.business-standard.com/india/news/india-achieves-241-mn-tonne-food-grain-production-in-fy11/141436/on](http://www.business-standard.com/india/news/india-achieves-241-mn-tonne-food-grain-production-in-fy11/141436/on); accessed on 29/02/2012). Our population is increasing day by day while our capacity to produce grains is not increasing in the same way, which suggests that we should enhance our production. It is believed

that nanotechnology increase the production and limit the disease attack and thus would be able to provide food to all.

Nanotechnology is a branch of science which deals with various aspects of research and technology. A joint committee of the British Royal Society and the Royal Academy of Engineering defined nanotechnology as the design, characterization, production, and application of structures, devices and systems by controlling shape and size at nanometer scale (Uskokovic, V, 2008). Nanoparticles are ultrafine particles whose size is greater than 1nm and smaller than 100 nm. There are many types of nanoparticles like Natural nanoparticle, incidental nanoparticles, quantum dots, organic polymers etc.

Nanotechnology deals with the matter at nanoscale (1-100 nm) dimensions. These materials

\* To whom all correspondence should be addressed.  
E-mail: sonica.dey@gmail.com

when reduced to the nanoscale show some properties which are different from what they exhibit on a macro scale, enabling unique applications. Nanoscience has brought revolution in different fields by helping develop processes and products that are hardly possible to evolve through conventional methods. The nanotechnology aided applications have the potential to change agricultural production by allowing better management and conservation of inputs of plant and animal production. A survey by Salamanca-Buentella et al. (2005) predicted several nanotechnology applications for agricultural production for developing countries within next 10 years. These included - (i) Nanoforms zeolites for slow release and efficient dosage of water and fertilizers for plants; drugs for livestock; nanocapsules and herbicide delivery (ii) Nanosensors for soil quality and for plant health monitoring; nanosensors for pest's detection (iii) Nanomagnets for removal of soil contaminants and (iv) Nanoparticles for new pesticides, insecticides, and insect repellents.

It is believed that nanotechnology would be able to manage food and nutritional security, that's why many large scale investments from global food corporations and many government societies from all over the world are investing in this branch (Roco, 2005; Sandler and Kay, 2006; Scrinis and Lyon, 2007). Crop productivity, soil health, disease management, water supply storage of food are the primary requirements for food storage, that can be full fill by nanotechnology (Kalpana Sastry et al., 2011). Some new Innovations are being made in agriculture sector, such as development of nanoseed within built pesticide effect. Nano encapsulation is also a nanotechnology based technique which has the potential to change nutritional composition, flavor etc., according to consumers requirement and physiological requirements. Presently nanotechnological intervention in agriculture focuses on three functional improvements (Observatory NANO, 2010).

1. Increased efficacy (with high solubility, stability and effectiveness)
2. Controlled release (in response to certain stimuli) and
3. Targeted delivery of fertilizers, plant growth regulators and biocides such as fungicides,

herbicides and pesticides.

Some nanotechnology based agriculture products that are under development age are:

- Nano- clay capsule containing biocontrol agents and growth enhancing enzymes that are designed for the slow release of active ingredients. Examples are pyrethroids such as y -cyhalothrin and ~-cypermethrin (Wang et al. 2007) and others like *Artemisia arborescens* L essential oil (Lai et al. 2006)...
- Nano-formulations (nanodispersions / nanoemulsions) of herbicide designed to attack the seed coating of weeds and prevent weed germination.
- Nanotechnology-enabled devices like stand-alone nano sensors linked into a GPS system for real-time monitoring of soil conditions and crop growth. Precision farming, with the help of smart sensors, will permit enhanced productivity in agriculture by providing accurate information, thus helping farmers and land managers in taking appropriate decisions.
- Nanoscale devices could be used for early disease detection and helping us to take preventive action. Nanotechnology will help in developing fertilizer and pesticide delivery systems which would be able to respond to environmental changes. Syngenta have obtained a patent for a nano pesticide called "gutbuster" microcapsules that contain pesticides engineered to break open in the alkaline conditions of an insect's stomach. Such nano-encapsulation techniques not only provide in-built pesticides for crops - in some ways similar to genetically modified Bt insecticidal crops -but also in-built switches to control the release and subsequent availability of pesticides (Scrinis and Lyons, 2007).
- By combining together Nanotechnology and Biotechnology a potent new tool would be produce that can manipulate genes and even produce new organism. For example, nanobiotechnologies enable nanoparticles, nanofibres and nanocapsules to carry foreign DNA and chemicals that modify genes (Torney ET al. 2007). In addition to the reengineering of existing plants, novel plant varieties may be developed using

synthetic biology; a new branch of technoscience that draws on the techniques of genetic engineering, nanotechnology and informatics (Lyons, 2010).

Nanotechnological Inventions will, therefore, enable development of novel agricultural chemicals, seeds, livestock, unknown plant growth promoting organisms, high-end use of agricultural water resources, high-precision farming systems etc., that will permit real time monitoring and management of farm operations.

#### **Agriculture and nanoparticles**

Numerous studies have reported the effects of nanoparticles on crops. But most of the experiments have been conducted in controlled conditions.

One study done by Khodakvskaya, 2009, found that carbon nanotubes enhance tomato seed germination and growth rate. CNTs were able to penetrate the thick seed coat and support water uptake inside seeds, a process which can enhance seed germination and growth of tomato seedlings. Lin and Xing 2007 found that suspension of MWCNTs have positive effect on the seed germination and root growth of radish (*Raphanus sativus*), rape (*Brassica napus*), rye grass (*Lolium perenne*), lettuce (*Lactuca sativa*), corn (*Zea mays*) and cucumber (*Cucumis sativus*).

TiO<sub>2</sub> NPs have been found to accelerating spinach growth (Zheng et al. 2005; Hong et al. 2005; Gao et al. 2006; Xuming et al. 2008). TiO<sub>2</sub> NP also improved plant growth by enhanced nitrogen metabolism (Yang ET al. 2006). They also found that TiO<sub>2</sub> nanoparticle have the capacity to increase nitrogen matabolism. In 2005 Watts and yang conducted an experiment on corn (*Zea mays*), cucumber (*Cucumis sativus*), soybean (*Glycine max*), carrot (*Daucus carota*) and cabbage (*Brassica oleracea*) with ALNPs and found that they reduce the elongation of roots of the plants and retard the plant growth. CNTs enhanced root elongation in onion and cucumber and nanotubes sheets were formed by both fCNTs and CNTs on cucumber root surface; however none entered into the roots. Cabbage and carrot were not affected by either form of CNTs. Root elongation in lettuce was inhibited with fCNTs and tomato was found to be most sensitive for CNTs with significant root length reduction. Reports by Nair et al. (2010). ZnO NPs showed no negative effects on seed

germination and root growth of zucchini seeds in hydroponic solution (Stampoulis et al., 2009) whereas the seed germination of rye grass and corn was inhibited by Zn NPs (35 nm) and ZnO NPs (15-25 nm) respectively (Lin and Xing, 2007).

Root growth of radish and rape incubated in Zn NP suspension was significantly inhibited (Stampoulis et al. 2009). They also found that ZnO NPs entered the root tissue of ryegrass which suggests that new nutrient delivery systems that exploit the nanoscale porous domains on plant surfaces can be developed (De Rosa et al., 2010).

In present time Nanoparticles made from Ag, Au, Si, Zn, Si etc are being tested as carrier for the delivery of pesticide, herbicide and fertilizers. Silicon nanoparticle is being tested for the delivery of avermectin and validamycin pesticides (Li et al., 2006; Liu et al 2006). Si nanoparticles have been reported to posse's insecticidal properties (Popat et al, 2012). Park et al, 2006 have found that when we apply composites of Si- Ag they found that it has potential to control various plant diseases.

#### **Effects of ENPs on microorganisms**

The main problem associated with the use of nanoparticles is their harmful effects which they leave after their applications. The exact toxic mechanism of ENPs is not completely known, the possible mechanisms through which they impart toxicology may include membranes disruption, proteins oxidation, genotoxicity, interruption of energy transfer pathway , reactive oxygen species (ROS) formation, toxin release etc. (Klaine et al.,2008).

Researches have found that ENPs like C<sub>60</sub> aggregates inhibited *Escherichia coli* and *Bacillus subtilis* (Fortner et al., 2005), and fullerene water suspensions (FWS) possess strong antibacterial activity and are more toxic to *B. subtilis* (Lyon et al., 2006). Metals like Ag, Au, Fe; Ti, Fe, Co-Zn-Fe etc have been found to be toxic against soil bacteria (Dinesh et al., 2012).

Silver nanoparticle was found to be toxic to *E. coli* and *Staphylococcus aureus* (Rai et al., 2009) and *B. subtilis* (Suresh et al., 2010). Silver nanoparticles biosynthesized by fungi possess potent activity against fungal and bacterial strains like *Aspergillus niger*, *Staphylococcus* sp., *E. coli* and *Bacillus* sp (Jaidev & Narasimha, 2010).

Water suspensions of nanosized TiO<sub>2</sub>, SiO<sub>2</sub> and ZnO were found to possess antibacterial

activity, antibacterial activity increases with particle concentration (Adams *et al* 2006). ZnO, Al<sub>2</sub> and TiO<sub>2</sub> NPs were found toxic to the nematode *Caenorhabditis elegans* inhibiting their growth especially the reproductive capacity (Wang *et al* 2008). Similarly, oxides of Zn, Cu and Ti NPs have been reported to be toxic to the micro algae *Pseudokirchneriella subcapitata* (Aruoja *et al*. 2009). Studies have shown that nanoparticles cause brain damage in fish and other aquatic species (Oberdorster, 2004; Oberdorster *et al*. 2007). Thus, from the studies conducted *in vitro* clearly shows that ENPs are hazardous to both micro-and macrofauna. In view of this problem researchers are trying to use these nanoparticles as carriers for delivery of the chemical fertilizers and pesticides to crops (De Rosa *et al*. 2010). It is important for us that we should set some standards for manufacturing, disposal and applications of these nanoparticles.

#### Environmental issues

Evidences suggests that nanoparticles have the ability to pose serious health, safety and environmental hazardous (Rocco and Bainbridge, 2001). ENPs which enter in our food either through crop application of nanoparticles, through packaging materials, as food additive have various nutritional, health and safety problems. Reports have indicated that ENPs can accumulate in plants and through plants they can find their way to humans, animals, microbial systems, aquatic system and soil (Nowack and Bucheli, 2007). Presently we have a very little knowledge about the impacts of nanoparticles on human health and environment but experiments conducted in laboratory have confirmed their toxicity against microorganisms. Still there is a lot to investigate about the health effects of foods that contain ENPs, or of workers handling nonmaterials.

The main problem associated with exposure of nanoparticles is that they may enter into the body through inhalation, digestion, skin etc. From there they found their way to circulatory system, into the lungs and may also cross the blood brain barrier (Scrinis 2006b).

In view of the growing importance of nanotechnologies the European Commission asked the independent experts of the Scientific Committee on Emerging and Newly Identified Health Risks (SCENIHR) to make scientific methodologies to access the potential hazardous

associated with nanotechnology applications. SCENIHR 2005 gave a report saying that the present methodologies require some modifications in order to deal with hazardous effects associated with nanotechnology. Equipments for routine measurements of free NPs in various media are not adequate. The available facilities to determine the fate of NPs in environment are not adequate. They also report that very little is known about physiology of nanoparticles

Presently, International Organization for Standardization (ISO) and the Organization for Economic Cooperation and Development (OECD) are seriously engaged in nanotechnology issues and have made a technical committee to develop international standards for nanotechnology (OECD, 2011). This technical committee, ISO/TC 229, is working to develop standards for terminology and nomenclature, metrology and instrumentation, including:

The OECD has established a Working Party on Manufactured Nanomaterials (WPMN) that is engaged in a variety of projects. The main working point of this agency includes:

- Database development on environmental health and safety (EHS) research
- To develop protocols for the manufacturing of nonmaterial's
- Testing of nonmaterial product manufactured to check whether it is safe or not.
- Develop a protocol regarding the manufacture and testing of nonmaterials.

EPA is actively working on this issue. The WPMN has made a list of manufactured nanoscale materials for environmental health and safety testing, including fullerenes (C 60), carbon nanotubes (single-and multi-walled), Ag-, Fe-, Ti-, Al-, Ce-, Zn-, Si-NPs, carbon black, polystyrene, dendrimers and nanoclays.(M. Anandarajn *et al.*, 2011).

#### CONCLUSION

Nanotechnology is an emerging branch of science with profound realized potential and outcome. It can be used as a powerful tool for the management of abiotic and biotic stresses, food and nutritional security, to increase yield of crop plants. However if not handled carefully the process and products formed by nanotechnology may pose

hazardous effects on health and environment. Hence we should maintain a strategy for using this science efficiently. Some government policies should also be formed to utilize this science. Controlled and wishful use of nanotechnology has a tremendous potential of improving agriculture sector.

### ACKNOWLEDGEMENTS

The authors are grateful for the financial support granted by the Indian Council of Agriculture Research (ICAR) Govt. of India under the Niche Area of Excellence on “Exploration and Exploitation of *Trichoderma* as an antagonist against soil borne pathogens” running in the Biocontrol Laboratory, Department of Plant Pathology, C.S.A. University of Agriculture and Technology, Kanpur, India.

### REFERENCES

- Observatory NANO. Nanotechnologies for nutrient and biocide delivery in agriculture production. Working Paper Version, 2010.
- Adams LK, Lyon DY and Alvarez PJJ. Comparative eco-toxicity of nanoscale  $TiO_2$ ,  $SiO_2$ , and  $ZnO$  water suspensions. *Water Res* 2006; **40**: 3527 -3532.
- Anandaraj M, Dinesh R, Srinivasan V and Harnza S. Nanotechnology in Agriculture: The Use of Novel Materials and Environmental Issues *The Botanica* 2011; 59 -61: 22 -34. 2011.
- Aruoja V, Dubourguier HC, Kasemets K and Kahru A. Toxicity of nanoparticles of  $CuO$ ,  $ZnO$  and  $TiO_2$  to microalgae *P. subcapitata*. *Sci Total Environ* 2009; **407**: 1461 -1468.
- Deendayal M . Mark E.Br, Debabrata M, Gobinda S, Priyabrata M, The use of microorganisms for the formation of metal nanoparticles and their application. *Appl Microbiol & Biotechnol*, 2006; **69**: 485-492.
- DeRosa Me, C Monreal, M Schnitzer, R Walsh and Y Sultan. Nanotechnology in fertilizers. *Nat Nanotechnol* 2010; **5**: 91.
- Dinesh R, Anandaraj M, Srinivasan V and Hamza S. Engineered nanoparticles in the soil and their potential implications to microbial activity. *Geoderma* 2012; 173 -174, 19 -27.
- Fortner JD, Lyon DY, Sayes CM, Boyd AM, Falkner JC, Hotze EM, Alemany LB, Tao YJ, Guo W, Ausman KD, Colvin VL and Hughes JB In C60 water: nanocrystal formation and microbial response. *Environ Sci Technol* 2005; **39**: 4307 -4316.
- Gao F. Mechanism of nano-anatase  $TiO_2$  on promoting photosynthetic carbon reaction of spinach: inducing complex of rubisco-rubisco activase. *Bioi Trace Elem Res* 2006; **111** : 239 -253.
- Hong F. Effect of nano- $TiO_2$  on photochemical reaction of chloroplasts of spinach. *Bioi Trace Elem Res* 2005; **105**: 269 -280.
- Jaidev LR and Narasimha G. Fungal media ted biosynthesis of silver nanoparticles, characterization and antimicrobial activity. *Colloids and Surfaces B: Biointerfaces* 2010; **81**: 430 -433.
- Kalpana Sastry R, HB Rashmi and NH Rao. Nanotechnology for enhancing food security in India. *Food Policy* 2011; **36**: 391 -400.
- Khodakovskaya M. Carbon nanotubes are able to penetrate plant seed coat and dramatically affect seed germination and plant growth. *ACS Nano* 2009; **3**: 3221 -3227.
- Kholoud MM, Abou El-Nour, Ala'a Eftaiha<sup>b</sup>, Abdulrhman Al-Warthan Synthesis and applications of silver nanoparticles. *Arabian Journal of Chemistry* 2010; **3**: 135- 140.
- Klaine SJ, Alvarez PJJ, Batley GE, Fernandes TF, Handy RD, Lyon DY, Mahendr S a, McLaughlin MJ and Lead JR. Nanomaterials in the environment: Behavior, fate, bioavailability, and effects. *Environ Toxicol Chern* 2008; **27**: 1825-1851.
- Lai F, Wissing SA, Muller RH and Fadda AM. *Artemisia arborescens* L essential oil loaded solid lipid nanoparticles for potential agricultural application: Preparation and characterization. *Aaps Pharrnscitech* 2006; 7.
- Li ZZ, SA Xu, Wen LX, Liu F, Liu AQ, Wang Q, Sun HY, Yu W and Chen JF. Controlled release of avermectin from porous hollow silica nanoparticles: Influence of shell thickness on loading efficiency, UV-shielding property and release. *J Controlled Release* 2006; **111**: 81 -88.
- Lin D and Xing B. Phytotoxicity of nanoparticles: inhibition of seed germination and root growth. *Environ Poll* 2007; **150**: 243 -250.
- Liu F, Wen LX, ZZ Li, Yu W, Sun HY and Chen JF. Porous hollow silica nanoparticles as controlled delivery system for water-soluble pesticide. *Mater Res Bull* 2006; **41**: 2268-2275.
- Lyon DY, Adams LK, Falkner JC and Alvarez PJ. Antibacterial activity of fullerene water suspensions: effects of preparation method and particle size. *Environ Sci Technol* 2006; **40**: 4360 -4366.
- Lyons K. Nanotechnology: Transforming food and the environment. *Food First Backgrounder* **16**: 1-4.
- Mahendra R, Alka Y, Aniket G. Silver nanoparticles: as a new generation of antimicrobials. *Biotechnology Advance*, 2009; **27**: 76- 83.

23. Meenal K, Shriwas A, Sharmin K, Voge W, Urban J, S.K.Kulkarni and K M Paknikar., Extracellular synthesis of silver nanoparticles by a silver-tolerant yeast strain MKY3. *Nanotechnology* 2003; **14**: 95-100.

24. Nair R, Varghese SH, Nair BG, Maekawa T, Yoshida Y and Sakthi D K. Nanoparticulate material delivery to plants. *Plant Sci* 2010; **179**: 154 -163.

25. Nelson D, Priscyla D M, Oswaldo L A, Gabriel IH De Souza, and Elisa E. Mechanistic aspects of biosynthesis of silver nanoparticles by several *F. oxysporum* strains. *J. Nanobiotechnol.* 2005; **3**: 1-7.

26. Nowack B Bucheli TD. Occurrence behavior and effects of nanoparticles in the environment. *Environ Poll* 2007; **150**: 5 -22.

27. Oberdorster E. Manufactured nanomaterials (fullerenes, C60) induce oxidative stress in the brain of juvenile largemouth bass. *Environ Health Perspect*, 2004; **112**: 1058 -1062.

28. Oberdorster G, Stone V and Donaldson K. Toxicology of nanoparticles: a historical perspective *Nanotoxicology* 2007; **1**: 2 -25.

29. Poovathinthodiyil R , Jie Fu , and Scott L. W. Completely “green” synthesis and stabilization of metal nanoparticles. *J Am Chem Soc* 2003; **125**: 13940-13941.

30. Popat A, Liu J, Hu Q, Kennedy M, Peters B, Lu QG and Qiao SZ. Adsorption and release of biocides with mesoporous silica nanoparticles. *Nanoscale* 2012; **4** : 970 -975.

31. Rai M, Yadav A and Gade A. Silver nanoparticles as a new generation of antimicrobials. *Biotechnol Adv* 2009; **27**: 76 -83.

32. Rajesh M and Dennis W. Metal Ion Resistance in Fungi: Molecular Mechanisms and their Regulated Expression. *J. Cell. Biochem*, 1991; **45**: 30-40.

33. Roco M. International perspective on government nanotechnology funding in 2005. *J Nailoparti Res* 2005; **7** : 707 -712.

34. Roco MC and Bainbridge WS. Societal implications of nanoscience and nanotechnology. viSET Workshop Report, 2001; 262.

35. Sandler Rand WD Kay. The National Nanotechnology Initiative and the Social Good. *The j Law, Med Ethics* 2006; **34**: 675 -681.

36. SCENIHR. Scientific committee on emerging and newly identified health risks. Opinion on the appropriateness of existing methodologies to assess the potential risks associated with engineered and adventitious products of nanotechnologies [<http://files.nanobio-raise.org/Downloads/scenihr.pdf> (accessed on 08/02/2012)], 2005.

37. Scrinis G. When Small isn't Beautiful. *Arena kfagazine* 2006b; **83**: 38 -40.

38. Scrinis G and K Lyons. The emerging nanocorporate paradigm: Nanotechnology and the transformation of Nature, food and agri-food systems. *International J Soc of Food & Agricul* 2007; **15**: 22 -44.

39. Stampoulis 0, Sinha SK and White JC. Assay-dependent phytotoxicity of nanoparticles to plants. *Environ Sci Technol* 2009; **43**: 9473-9.

40. Suresh AK, Pelletier0, Wang W, Moon JW, Gu B, Mortensen NP, Allison OP, Joy DC, Phelps TJ and Ooktycz MJ. Silver nanocrystallites: Biofabrication using *Shewanella oneidensis*, and an evaluation of their comparative toxicity on gram-negative and grampositive bacteria. *Environ Sci Technol* 2010; **44**: 5210 -5215.

41. T Prameela D, S Kulanthaivel, Deeba K, Jyoti LB, N Prabhakaran and N Srinivasa. Biosynthesis of silver nanoparticles from *Trichoderma* species. *Indian J. Exp. Biol.* 2013, **51**: 543-547.

42. Torney F, Trewyn B, Lin V and Wang K. Mesoporous silica nanoparticles deliver DNA and chemicals into Plants. *Nat Nanotechnol* 2007; **2**: 295-300

43. Vuk U. Nanomaterials and Nanotechnologies: Approaching the crest of this big wave. *Current Nanoscience*, 2008; **4**: 119.

44. Wang H, Wick RL and Xing B. Toxicity of nanoparticulate and bulk ZnO, Al<sub>2</sub>O<sub>3</sub> and TiO<sub>2</sub> to the nematode *Caenorhabditis elegans*. *Environ Poll* 2008; **157**: 1171 -1177.

45. Wang L Z Li G Zhang J, Dong and Eastoe J. Oil-in-water nanoemulsions for pesticide formulations. *J Colloid Interf Sci* 2007; **314**: 230 -235.

46. Xuming W). Effects of nano-anatase on ribulose-I, 5-biphosphate carboxylase/oxygenase mRNA expression in spinach. *Biol Trace Elem Res* 2008; **126**: 280 -289.

47. Yang F. Influences of nano-anatase Zheng L, F Hong, S Lu and C Liu. Effects of TiO<sub>2</sub> on the nitrogen metabolism of growing spinach. nano-TiO<sub>2</sub> on strength of naturally aged seeds. *Bioi Trace Elem Res* 2005; **110**: 179 -190.

48. Yang L, Watts DJ. phytotoxicity of alumina nanoparticles. *Toxicology - Letters* 2005; **158**: 122 -132.

49. Zheng L, Hong F, Lu S and Liu C Effects of TiO<sub>2</sub> on the strength of naturally aged seeds and. growth of spinach. *Bioi Trace Elem Res* 2005; **104**: 83 -92.

50. Zhou X, A Striolo and PT Cummings. C60 characteristics may play an important role in binds to and deforms nucleotides Particle surface. *Biophys J* 2005; **89**: 3856- 3862 .

© The Author(s) 2016. **Open Access.** This article is distributed under the terms of the [Creative Commons Attribution 4.0 International License](http://creativecommons.org/licenses/by/4.0/) which permits unrestricted use, sharing, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.