

The Effect of Cytokinin and Iron on Biological Nitrogen Fixation in Chickpea Cultivars under Dryland Conditions

Kianoush Hamidian^{1*}, Ahmad Naderi^{2*}, Shahram Lak³ and Islam Majidi⁴

¹Department of agronomy, Callege of Agriculture,

Khuzestan Science and research Branch, Islamic Azad University, Ahvaz, Iran.

²Faculty of Agriculture and Natural Resource Research Center of Khuzestan, Ahvaz, Iran

³Department of agronomy, Callege of Agriculture, Ahvaz Branch, Islamic Azad University, Ahvaz, Iran.

⁴Department of Plant Breeding, Callege of Agriculture, Tehran Science and Research Branch, Islamic Azad University, Tehran, Iran.

<https://doi.org/10.22207/JPAM.10.1.51>

(Received: 02 September 2015; accepted: 17 January 2016)

In order to evaluate the effect of cytokinin and iron on biological nitrogen fixation in chickpea cultivars, experiment was conducted as a factorial experiment in randomized complete block design in four replications. The treatments consisted of three cultivars: (Hashem, Azad and local mass), four levels of iron [control (lack of consumption), 5, 10 and 15 ppm] and four levels of cytokinin (6-Benzyl aminopurin) [control (lack of consumption), 5, 10 and 15 ppm]. Local mass with 3.69, 6.1 and 3.78 percent, had the highest nitrogen percent in the above ground, noude and grain. By increasing concentrations of iron, increased nitrogen percent in aboveground and grain, but in noude increased up to 10 ppm and then decreased. However, by increasing concentrations of cytokinin, the nitrogen percent of grain increased, but nitrogen percent of aboveground and noude increased up to 10 ppm and then decreased. Local mass with 129.8 kg/ha had the highest nitrogen fixation. Effects of iron and cytokinin on nitrogen fixation was positive, and by increase their concentration, nitrogen fixation increased and in concentration of 15 ppm content of fixed nitrogen were 130.9 and 129.8kg/ha respectively. From of all fixed nitrogen in plant, Contribution of aboveground, grain and noude were 46.19, 46.41, and 7.4 percent.

Keywords: Chickpea, biological nitrogen fixation, cytokinin, iron, aboveground, noude and grain.

Nitrogen is the main nutrient in chemical compounds, metabolism and protein synthesis^{18,25}. Although the plant nitrogen is about 1 to 2 and the maximum 4 to 6 percent, however, nitrogen is the limiting factor for crop production in the world²⁵. Biological nitrogen fixation in legumes can be done effectively, therefore, plant can grow by consuming less of nitrogen fertilizer¹³. The importance of biological nitrogen fixation are to replace the nitrogen lost by culture, leaching, volatilization, runoff, erosion and denitrification⁹. Chickpea is an annual plant that by providing the

environmental conditions, through symbiosis with Rhizobium bacteria causing nitrogen fixation and don't need to nitrogen fertilizer and in desired rotations, will provide the part of required nitrogen for next crop¹⁰. The amount of nitrogen fixation in chickpea are different from zero to 248 kg/ha in the growing season^{17,25}. Islam *et al.*,¹⁵ estimated the amount of nitrogen fixation 52 to 55 percent from of all nitrogen plant. Evers *et al.*,⁹ in a study on clover cultivars, estimated up to 296 kg/ha nitrogen fixation.

Cytokinins, generally adenine derivatives, are plant hormones that promote growth and cell division and essential for plant growth and development^{4,11} and reducing their amount in the root causes limiting of protein synthesis and reducing of legheamoglobin, restriction of node

* To whom all correspondence should be addressed.
Tel/Fax: 986633220112;
E-mail: keanoush.hamidian@gmail.com

respiration, change in node metabolism or effect on nitrogenase enzyme activity and finally reduced nitrogen fixation¹². Cytokinins play an important role in forming of plant mutualism with bacteria and mycorrhiza through symbiont. The hormone to stimulating of symbiosis, increase activity of nodulation genes and facilitate transfer of nutrients between plant and symbiotic²⁴.

Iron is a key factor to activity of many enzymes and often is very important for catabolism activity. The low solubility of iron in water, limit biochemical processes such as photosynthesis and nitrogen fixation²².

genotype like environment and growth period are important factors that affect on plant growth^{16,18}. Species and genotypes differ in susceptibility to iron deficiency and traits that affects by genotypes ability on the use of iron, are dry matter and grain yield²⁸. Mansour *et al.*,¹ showed that there was significant difference between genotypes of chickpea in nitrogen uptake.

MATERIALS AND METHODS

This study was performed at Grit Agricultural Services Center (48° 18' long. E; 33° 32' lat. N, and 1175 m above sea level) located in 35 kilometers Eastern-North of Khorramabad of lorestan province in 2011 and 2012 .According to meteorological long term, mean annual precipitation is 425 mm.

Experiment was conducted as a factorial experiment in randomized complete block design in four replications. The treatments consisted of three cultivars: (Hashem, Azad and and local mass), four levels of iron [control (lack of consumption), 5, 10 and 15 ppm] and four levels of cytokinin (6-Benzyl aminopurin) [control (lack of consumption), 5, 10 and 15 ppm]. Each plot consisted of five rows with 6 m length that space between rows and plants space on rows were considered 25 and 15 cm, respectively. Planting was done according to current planting date in area and atmospheric conditions in March 23, 2011 and March 13 , 2012.

Before planting, soil sample prepared from the 0-30 cm depth and were measured the physical and chemical properties of soil (Table 1). In this study, neither macro and nor micro fertilizers were applied. Iron($\text{FeSo}_4 \cdot 7\text{H}_2\text{O}$) was sprayed in

3-4 leaf stage(45 DAP¹) and cytokinin(6-Benzil aminopurine) sprayed 10 days later.

Traits and their measurements

At 50% flowering stage, from each plot, were eliminated two lateral rows and a half meters from start of middle rows as border effects. from three middle rows, selected three plants with 30 cm³ of soil their around and after washing the soil around the roots, the nodules on the roots were collected, then, were dried for 72 hours in air and weighed with a digital scale with an accuracy of 0.001. At this stage, from each plot, cut six plants and was prepared composite sample. as well as, in harvest stage, 20 seeds selected randomly from each plot. in laboratory were measured nitrogen percent of nodes, aboveground and grain by Kjeldahl method.

Nitrogen fixation

Based on nitrogen percent of aboveground, noude and grain, fixed Nitrogen in any part of plant and total nitrogen fixation in plant were calculated by use of following formula:

$$\begin{aligned} NF &= DW \times \%N \\ NF_g &= DW_g \times \%N_g \\ NF_n &= DW_n \times \%N_n \\ TNF &= NF_g + NF_s + NF_n \end{aligned}$$

that:

NF : Nitrogen fixation in grain(kg/ha)

NF_n : Nitrogen fixation in nodule (kg/ha)

NF_a : Nitrogen fixation in above ground plant(kg/ha)

TNF Total nitrogen fixation in plant(kg/ha)

DW: Dry weight of plants(kg/ha)

N%: Nitrogen percen.

Statistical analysis

Data were combined analyzed using MSTAT-C software and graphs were drawn using EXCEL2010. Means were compared using Duncan's Multiple Range Test.

RESULTS AND DISCUSSION

Dry weight of aboveground, grain and node

The results showed that local mass with 8.164, 6.15 and 7.156 gr/m² had the highest dry weight of aboveground, grain and node, respectively. (Table 2). This result were confirmed by James *et al.*,¹⁵ Seyfi²⁸, Biabani and Carpenter⁵ and Khan *et al.*,¹⁷.

Genotype like environment and growth period are important factors that affect plant growth^{16,18}. James *et al.*,¹⁶ showed that dry weight of aboveground was significantly affected by cultivar. In a study by Khazaei *et al.*,¹⁹, genotypes in respect of noude number, noude weight and dry weight of aboveground were significantly different, Mcc480 genotype had more number of nodes with higher specific weight to Mcc876 genotype.

With increasing iron concentration up to 10 ppm, dry weight of aboveground and nodes

increased then decreased, while with increasing concentration of iron, grain dry weight had increased (Table 3).

Species and genotypes differ in susceptibility to iron deficiency and traits that affects by genotypes ability on the use of iron, are dry matter and grain yield²⁸. In study by Khan *et al.*,¹⁷, response of chickpea cultivars to iron use was positive and obtained highest dry weight of shoot and grain by spray of 2 kg/ha ferrous sulfate. Saifi²⁸ to study of iron on grain yield of chickpea

Table 1. Physical and chemical properties of soil test site

Feature year	%N	P(ppm)	K(ppm)	Fe(ppm)	Mn(ppm)	Zn(ppm)	S(ppm)	%humus	Texture	pH
2010-2011	0.085	14.8	342.2	8.88	8.68	1.58	80.62	1.1	Silty loam	7.6
2011-2012	0.069	11.4	367.3	7.42	6.65	0.95	92.45	0.9	Silty loam	7.3

Table 2. The effect of cultivar on dry weight, nitrogen percent and fixed nitrogen in aboveground, grain and node

Cultivar	Dry weight (g / m ²)			Nitrogen percent			Ffixed nitrogen (kg / ha)			Total fixed nitrogen (kg/ha)
	aboveground	grain	node	aboveground	grain	node	aboveground	grain	node	
Hashem	161.5b	152.1b	12.6b	3.63c	3.69b	6.06b	58.6b	56.2b	7.6b	122.4b
Azad	161.1b	152.0b	12.1c	3.65b	3.65b	6.05b	58.7b	55.6b	7.3c	121.7b
Local mass	164.8a	156.7a	15.6a	3.69a	3.78a	6.10a	60.8a	59.3a	9.5	129.8a

Table 3. The effect of iron on dry weight, nitrogen percent and fixed nitrogen in aboveground, grain and node

Iron ppm	Dry weight (g / m ²)			Nitrogen percent			Ffixed nitrogen (kg / ha)			Total fixed nitrogen (kg/ha)
	aboveground	grain	node	aboveground	grain	node	aboveground	grain	node	
0	160.7c	139.1c	12.4d	3.64b	3.69b	5.91d	58.5b	51.3c	7.4d	117.2d
5	161.2b	153.4b	13.5c	3.64b	3.7b	6.08c	58.7b	56.7b	8.2c	123.6c
10	164.3a	155.8b	14.1a	3.67a	3.7b	6.18a	60.3a	57.6b	8.7a	126.6b
15	164.4a	166.2a	13.7b	3.67a	3.76a	6.15b	59.9a	62.5a	8.4b	130.8a

Table 4. The effect of cytokinin on dry weight, nitrogen percent and fixed nitrogen in aboveground, grain and node

Cytokinin ppm	Dry weight (g / m ²)			Nitrogen percent			Ffixed nitrogen (kg / ha)			Total fixed nitrogen (kg/ha)
	aboveground	grain	node	aboveground	grain	node	aboveground	grain	node	
0	160.1d	140.6d	12.7c	3.56c	3.61c	5.88d	56.9c	50.7d	7.5d	115.1d
5	161.7c	149.6c	13.4b	3.64b	3.69b	6.00c	58.8b	55.2c	8.0c	122.0c
10	163.2b	159.4b	13.8a	3.71a	3.74a	6.24a	60.5a	59.7b	8.6a	128.8b
15	164.9a	164.8a	13.8a	3.70a	3.79a	6.19b	61.1a	62.5a	8.5b	132.1a

cultivars, reported that iron were effective on grain yield and Arman cultivar in 15 kg/ha iron treatment had highest yield (692.56 kg / ha)

Grain weight is one of the important yield components that will be influenced by genetic characteristics such as grain potential, grains competition, grain filling period and environmental conditions²⁹. Amini dehogy and dadkhah² reported that iron caused significantly different in number and weight of nudes. Welch³⁴ stated that iron with participation in metabolism of hydrocarbon material and protein and their transfer, and effects on reproductive process, had significant role in increasing the number and weight of grains. Rai *et al.*²⁶ reported that iron deficiency reduced number and weight of noudes in chickpea.

With increasing concentrations of cytokinin, dry weight of aboveground, seed and node significantly increased and highest dry weight was obtained in 15 ppm (Table 4). Hormone use has a great impact in achieving high-yield. Cytokinins are plant hormones that as regulate growth and development has an important role in preventing the aging of organs and nutrient mobility³. Zulkarnyan *et al.*³⁶ studied the effect of cytokinin on rice, the results showed that use of cytokinin increased the number of tillers, spikes and finally yield. Najel *et al.*²³ examined the effects of cytokinin on soybean yield, results showed that consumption $5 \times 10^{-6} \mu$ increased aboveground dry weight, number and weight of seeds. In this study, the doses more than $5 \times 10^{-6} \mu$ decreased 30% number of grains and 39% grain weight compared with control. In this study, the reason of produce more grain, effect of cytokinin in reducing stimulate production of flowers and prevent of flower abortion were expressed. Fatima *et al.*¹⁰ examined effects of IAA, ABA and kinetin on growth, yield and yield components of chickpea in spray and seed soaking in solution of growth hormones, results indicated an increase in aboveground dry weight in both spray and seed soaking by Kinetin, and increasing grain weight by seed soaking in Kinetin, the maximum grain weight in kinetin were attributed to transfer of assimilates from vegetative parts to reproductive parts. Based on research conducted by Iqbal *et al.*¹⁴, cytokinin had a positive effect on seed size, this is due to the effect of cytokinin on the processes of accumulation of nutrients in the grain, continuation of grain filling period and increase transfer nutrients from roots

to grains. kinetin as growth-stimulating hormone increases the activity of nitrogenase enzyme in chickpea root nodes⁷, and by increasing the amount of legheamoglobin influences efficiency and viability of the nodes³³. Fatima *et al.*¹⁰ examined effects of IAA, ABA and Kinetin and two strains of Rhizobium on growth, yield and nitrogen fixation in chickpea under natural conditions, the results showed that maximum weight of nodes obtained in seeds treated with kinetin and Tal 620 rhizobium race.

Nitrogen percent of aboveground, node and grain

Local mass had highest percent of nitrogen in aboveground, node and grain (Table 2). N concentration in plant depends on factors such as amount of nitrogen in soil, plant type, plant organs and plant growth stage²⁰. Mansour *et al.*²¹ showed that there was significant difference between genotypes of chickpea in terms of nitrogen uptake.

With increasing concentrations of iron and cytokinin up to 10 ppm, increased nitrogen percent of aboveground and node, while the highest percent of grain nitrogen obtained in 15 ppm (Tables 3 and 4). Singh *et al.*³¹ to study of effect of iron concentration on chickpea found that with increasing iron concentration, content and nitrogen uptake were significantly increased, in this study, iron caused more nitrogen absorption and transfer it between different parts of plant. the positive reaction of nitrogen percent to iron is due to its undeniable role in symbiotic bacteria and plant roots² and Iron involvement in the building of enzymes such as nitrogenase, nitrate and nitrite reductase and legheamoglobin³². Singh *et al.*³⁰ found a positive correlation between nitrogen and cytokinin in the plant species.

Nitrogen fixation

The results showed that the effects of iron and cytokinin on the amount of nitrogen fixation in different parts of plant and total nitrogen fixation were significant. Local mass with 60.87 kg/ha had the highest content of nitrogen fixation, while Hashem and Azad with 58.71 and 58.59 kg/ha were analyzed in a group, respectively (Table 2). Nitrogen fixation by chickpea is determined by two factors:

1. Amount of nitrogen accumulation in plant during growth period
2. Amount of nitrogen absorbed by plants from of all fixed nitrogen⁸.

Biabani *et al.*,⁵ studied nitrogen fixation of 40 cultivars that were collected from Iran, Afghanistan, Iraq, Ethiopia, Mexico, India, Pakistan, Turkey and United States. They reported significant differences between cultivars and Genotype 254549 from Iraq to 0/084 g/plant had the highest nitrogen fixation.

The effect of iron and cytokinin were significant on content of fixed nitrogen in aboveground, nodes and grain and total fixed nitrogen. Except node, with increasing iron and cytokinin concentrations, increased amount of nitrogen fixation (Tables 3 and 4). Khan *et al.*,¹⁷ reported that iron and cultivar were affected on nitrogen fixation and most fixed nitrogen obtained by karek cultivar (Kabuli type) that treated with 2 lit/ha iron. Calsikan *et al.*,⁶ reported that iron deficiency had negative effect on nitrogenase enzyme activity and nitrogen fixation in soybean. Amini Dehoghi and Dadkhah² by investigation effects of iron and zinc fertilizers on nitrogen fixation found that in salinity condition, iron fertilizer moderated salinity effects and increased nitrogen fixation.

Fatima *et al.*,¹⁰ to survey of effect of growth regulators on nitrogen fixation in chickpea found that kinetin caused an increase in total fixed nitrogen. Kinetin as growth-stimulating, increased activity of nitrogenase enzyme in nodes⁷. Rao²⁷ reported irritant effect of kinetin on nitrogen fixation efficiency. Overall, local mass in concentrations of 15 ppm iron and cytokinin with 142.1 kg/ha had highest nitrogen fixation. Of all fixed nitrogen in plant, the share of aboveground, grain and node were 46.19, 46.41 and 7.4 percent, respectively.

CONCLUSION

Cultivars had different responses to nitrogen fixation in different parts of plant. It seems that local mass due to the long term adaptation to environmental conditions and possibility of better strains of Rhizobium bacteria in soil, had been able to fix a higher content of nitrogen. Cytokinin as growth-stimulating hormone, and iron as an essential component of leghemoglobin, increased nitrogenase enzyme activity in root nodes and caused more nitrogen fixation efficiency. It is recommended to increase fixed nitrogen, increases

harvest index, till to increasing the grain yield, increases total amount of fixed nitrogen.

REFERENCES

1. Ali. S. and A, Bano. Leaf and nodule senescence in chickpea (*Cicer aritinum L*) and the role of plant growth regulators. *Pak.J.* 2008; **40**(60): 2481-2492.
2. AminiDehoghi M, Dadkhah A., Effect of iron and zinc fertilizers on growth, nodulation and nitrogen fixation in chickpea under saline conditions. 4th Conference of Pulses. *Arak.* 2011.
3. Argueso CT, Ferreira FJ, Kieber JJ, Environmental perception avenues: the interaction of cytokinin and environmental response pathways. *Plant, Cell and Environment.* 2009; **32**: 1147–1160.
4. Arteca RN., Plant Growth Substances: Principles and applications. Jahad-e-daneshgahi Press. Mashhad, Iran, 2000.
5. Biabani A, Carpenter-Boggs L, Clarice J, Taylor L, Jeffre., Nitrogen fixation Potential in global chickpea mini-core collection. *Biol Fertil Soils* 2011; **47**: 679-685.
6. Calsikan S, Ozkaya I, Calsikan ME, Arslan A., The effects of nitrogen and iron fertilization on growth, yield and fertilizer use efficiency of soybean in Mediterranean type soil. *Field Crops Research.* 2008; **108**(2): 126-132.
7. Dayal J, Baharti S., Effects of Kinetin, cycocel and colchicine on nitrogenous activity and national symposium on growth and differentiation in plant. (Eds.): K.K. Dhir, I.S, 1991.
8. Doughton JA, Vallis L, Saffigna PG., Nitrogen fixation in chickpea. 1. Influence of prior cropping or fallow, nitrogen fertilizer and tillage. *Australian Journal of Agricultural Research* 1993; **44**: 1403-13.
9. Evers GW, Parsons MJ., Estimated N₂-Fixation of cool-season annual clovers by the difference method. *Crop science* 2011; **51**(5): 2276-2283.
10. Fatima Z, Aslam M, Bano A., Chickpea nitrogen fixation increases production of subsequent wheat in rainfed system. *Pak J Bot* 2008; **40**(1): 369-376.
11. Fransworth E., Hormones and shifting ecology throughout plant development. *Ecology*, 2004; **85**(1): 5-15.
12. Goicoechea N, Antolin, MC, Strand M, Sanchez-Diaz M., Root cytokinins, acid phosphatase and nodule activity in drought- stressed mycorrhizal or nitrogen-fixing alfalfa plants. *Journal of Experimental Botany* 1996; **47**(298): 683-686.
13. Hardarson G, Danso SKA., Methods for

- measuring biological nitrogen fixation in grain legumes. *Plant and soil*. 1993; **152**: 19-23.
14. Iqbal N, Basrash MA, Rehman KH., Evaluation of vigour and oil quality in cottonseed during accelerated ageing. *Int J Agri Biol* 2002; **4**: 318-322.
 15. Islam M, Saleem M, Safdar A, Khlid R, Hassan F, Mahmood A and Subhani A., Growth, Nitrogen Fixation and Nutrient Uptake by Chickpea (*Cicer arietinum*) in Response to Phosphorus and Sulfur Application under Rainfed Conditions in Pakistan. *J Agric Biol* 2011; **13**(5):725-730
 16. James E, Mackown TC, Rao SC., Dry weight and nitrogen content of chickpea and winter wheat grown in pots for three rotations. *Journal of Plant Nutrition* 2007; **30**: 1541-1553.
 17. Khan N, Tariq M, Ullah M, Muhammad D, Khan I, Rahatullah K, Ahmed N, Ahmed S., The Effect of Molybdenum and Iron on Nodulation, Nitrogen Fixation and Yield of Chickpea Genotypes (*Cicer Arietinum L*). *IOSR-JAVS*. 2014; **7**(1): 63-79.
 18. Khattak SG, Khan DF, Shah SH, Madani MS, Khan T., Role of rhizobial inoculation in the production of chickpea crop. *Soil and Environ*. 2006; **25**(2): 143-145.
 19. Khazaei H, Parsa M, HosseinPanahi M., Effects of Rhizobium inoculation on nodulation of Desi and Kabuli varieties in soil moisture regimes. *Journal of iran agricultural research*. 2008; **6**(1): 97-89.
 20. Malakouti M, Homaei J, Fertility of soils in arid and semi-arid in Iran (problems and solutions) second edn, Tarbiat Modarres University. Tehran, 2004.
 21. Mansur CP, Palled YB, Halikatti SL, Salimath PM, Chetti MB, Uptake of nitrogen phosphorus and protein content in Kabuli chickpea genotypes as influenced by plant densities and phosphorus levels. *J Agric Sci* 2009; **22**(1): 5-7.
 22. Mora V, Bacaicoa E, Zamarreno AM, Aguirre E, Garnica M, Fuentes M, Garcia-Mina JM) Action of humic acid on promotion of cucumber shoot growth involves nitrate-related changes associated with the root-to-shoot distribution of cytokinins, polyamines and mineral nutrients. *J Plant Physiol* 2010; **167**: 633-642.
 23. Nagel L, Brewster R, Riedell WE, Reese RN, Cytokine Regulation of flower and set in soybeans (*Glycine max L.*) *Annals of Botany* 2001; **88**:27-30.
 24. Ng PP, Cole AL, Jameson PA, McWha JA, Cytokinin production by ectomycorrhizal fungi. *New Phytologist*. 1982; **91**: 57-62.
 25. Parsa, M., Bagheri, A, Pulses. University of Mashhad Publication. Mashhad, 2008.
 26. Rai R, Singh SN, Prasad V., Effect of pressmud amend pyrite on symbiotic N₂-fixation, active iron contents of nodules, grain yield and quality of chickpea (*Cicer arietinum L*) genotypes in calcareous soil. *Journal of Plant Nutrition* 1982; **5**: 905-913.
 27. Rao LVM, Neeraj M, Datta M, Sudhir K, Influence of cytokinin and phytochrome on nitrate reductase activity in etiolated leaves of maize. *Photochemistry* 1984; **23**:1875-1884.
 28. Reddy KR, Delaune RD, Ebusek WFD, Koch MS, Long-term nutrient accumulation rates in the Ever-glades. *Soil Sci Soc Am J* 1993; **57**:1147-1155.
 28. Seifi A, Khazaei A, Rafie, M., Effect of iron on yield and yield components of chickpea cultivars under rainfed conditions. MS thesis. Islamic Azad University. Khorramabad, 2008.
 29. Shanahan JF, Smith DH, Welsh JR., An analysis of post-anthesis sink-limited wheat grain under various environments. *Agronomy Journal* 1984; **76**: 611-615.
 30. Singh B, Narang MP., A comparison of chemical composition, cell-wall content, digestibility and degradation kinetic characteristics as predictors of forage intake. *Indian J Anim Sci* 1992; **62**: 369-373.
 31. Singh M, Chaudhary Sharma SR, Rather MS, Effect of same micronutrients on content and uptake by chickpea (*cicer arietinum L*). *Agri SCI Digest* 2004; **24**(4):268-270.
 32. Vanoni MA, Curti B, Structure-function studies on the iron-sulfur flavin-enzymes glutamate synthase: an unexpectedly complex self-regulated enzyme. *Arch Biochem Biophys*. 2005; **433**: 193-211.
 33. Wanger GH, Beck E, Bernd M, Cytokinins in the perennial herb *Urtica dioica L.* as influenced by its nitrogen status. *Planta*, 1993; **190**: 511-518.
 34. Welch RM., Micronutrient nutrition of plants. *Critical reviews in plant sciences*. 1995; **14**:49-82.
 35. Ziaian A, Malakouti M J., Evaluation the greenhouse effect of Fe, Mn, Zn and Cu on wheat production in Fars province soil and water *Journal*. 1999; **12**(6): 24-32.
 36. Zulkarnain WM, Ismail MR, Saud HM, Othman R, Habib SH, Kausar H, Effect of synthetic cytokine precursor on growth and yield of chickpea under limited water. *Journal of food, Agriculture & environmental* 2013; **11**(2):372-375.

© The Author(s) 2016. Open Access. This article is distributed under the terms of the [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) which permits unrestricted use, sharing, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.