

Antimicrobial Effectiveness of $[\text{Cu}(\text{pzdc})(\text{phen})_2] \cdot 5.5\text{H}_2\text{O}$ Complex on Some Microorganisms

Cihan Darcan^{1*}, Özge Kaygusuz² and Tülin Korkmaz²

¹Department of Molecular Biology and Genetics, Faculty of Arts and Sciences,
Bilecik Seyh Edebali University, Bilecik-Turkey.

²Department of Molecular Biology and Genetics, Institute of Sciences,
Bilecik Seyh Edebali University, Bilecik-Turkey.

(Received: 24 March 2013; accepted: 08 May 2013)

This study was conducted to find new chemical substances that are highly effective on selected microorganisms from among eight Cu(II) pyrazine-2,3-dicarboxylate complexes, which have been previously synthesized and characterized. The MIC value of the $[\text{Cu}(\text{pzdc})(\text{phen})_2] \cdot 5.5\text{H}_2\text{O}$ complex (8) was determined to be in the range of 15-40 $\mu\text{g/ml}$ for all bacteria (except the clinical isolate *Pseudomonas aeruginosa*). However, the minimum microbiocidal concentrations (MMC) (30-200 $\mu\text{g/ml}$) were determined to be much higher than the MIC values. In time-kill assays, all the bacteria were fully finished after 120 minutes at 18 $\mu\text{g/ml}$ and after 240 minutes at 4.5 $\mu\text{g/ml}$ from 5.54 log, the initial number of bacteria at 37 °C in the lake water. However, all the bacteria were not fully inhibited even at 18 $\mu\text{g/ml}$ at 10 °C. The same effect was not seen on *Candida albicans* in spite of the low MIC value (25 $\mu\text{g/ml}$). $[\text{Cu}(\text{pzdc})(\text{phen})_2] \cdot 5.5\text{H}_2\text{O}$ complex has broad-spectrum biostatic activity against microorganisms at the MIC values. The mechanism of action is not yet clear.

Key words: Antimicrobial activity, Disinfectant, Pyrazine-2,3-dicarboxylate complex.

A significant percentage of clinical isolates, including species of methicillin-resistant *Staphylococcus aureus* (MRSA), *Pseudomonas aeruginosa*, *Streptococcus pneumoniae*, *Enterococcus faecalis*, *Acinetobacter baumannii* and *Morganella morganii* are resistant to commonly used antibiotics, such as new-lactams, glycopeptides, fluoroquinolones, linezolid and macrolides¹⁻⁶. At the same time, these bacteria generates serious public health problems which are called nosocomial infections because of the infection and superinfection of hospitalized patients. Especially, the situation is critical in the

case of treatment of hospital infections caused by *S. aureus* and *P. aeruginosa*. Every year, two million hospital infections occur in the United States, and 90.000 people are reported dead⁷. The annual prevalence of hospital infections in a report including the nine countries has been reported to be 4-10%. In Turkey, this ratio varies between 1% and 8.6%. The frequency of hospital acquired infection in intensive care units is reported to be 50%, with the death ratio being 16%⁷. The emergence of antibiotic resistance among pathogenic bacteria has led to major research efforts to find alternative antimicrobial therapeutics. The spread of resistant bacteria causes problems of infection control in society. Clearly, new antimicrobial agents that can be effective at low doses, with broad-spectrum or species-specific activity, are required to combat

* To whom all correspondence should be addressed.
Tel: +90 228 2141161-1174; Fax: +90 228 2141162;
E-mail: cihandarcancan@yahoo.com

an ever-increasing resistance problem among microorganisms. New antimicrobial agents are required for not only human diseases but also plant and animals diseases, for disinfection or sterilization of environment, water and wastewater.

In recent studies on the production of antimicrobial agents, metal ions such as Cd(II), Pd(II), Ag(I) and Au(I) and biocations such as Co(II), Ni(II), Cu(II), Fe(III), Cr(III), Mn(III) and Zn(II) were shown possess high activity to inhibit the development of bacterial resistance^{8,9}. Positively charged metal centers are preferred in the production of antimicrobial drugs because they facilitate interaction with negatively charged biomolecules such as nucleic acids, ATP and protein components. Metal ions exert their effect by exchange of their metals or by disturbing to the internal and external coordination of active region to the structural integrity of enzymes. Metal ions allows longer time interaction through strong covalent or ionic bonds with the target molecules, compared to organic antimicrobial agents¹⁰.

Although antimicrobial agents containing metals have high activity, some of them can not be used as antimicrobials, disinfectants, antiseptics due to their toxicity, drug resistance, and cellular metabolism of the complexes. Among antimicrobial agents, only silver complexes can be used to prevent bacterial infections in the treatment of burns, and to prevent the development of ophthalmia neonatorum^{11,12}. Silver sulfadiazine (AgSD), a sulfa drug, is used to prevent and treat infections of second- and third-degree burns. Silver is an effective antimicrobial agent with low toxicity and hence is important especially in the treatment of burn wounds^{11,13}.

The purpose of this study is to find new chemicals that can be used in the war against microorganisms. For this purpose, complexes that were synthesized and characterized by Yesilel *et al.*^{14, 15, 16, 17} were selected, and the antimicrobial effects of these complexes were investigated.

MATERIALS AND METHODS

Microorganisms and metal complexes used in this study

The bacteria (seven wild type and six clinical isolates) and yeast used in this study are listed in Table 1. The clinical isolates were collected

from the Faculty of Medicine, University of Ondokuz Mayıs. All the clinical isolates were isolated from patients in the hospital. The eight pyrazine-2,3-dicarboxylate complexes, the pyrazine-2,3-dicarboxylic acid (H_2pzdc) and phenanthroline (phen) ligands used in this study are shown in Table 2. The preparation of the metal complexes, and their characterization steps have been previously presented in articles published by Yesilel *et al.*^{14, 15, 16, 17}. The $Cu(pzdc)(phen)_2 \cdot 5.5H_2O$ complex is shown at Fig. 1¹⁵.

Minimum Inhibitory Concentration (MIC) tests of complexes

The stock solutions of metal complexes were prepared with distilled water (30 mg/ml). The in-vitro susceptibility tests (MIC tests) were determined by broth microdilution carried out in accordance with the guidelines of the National Committee for Clinical Laboratory Standards¹⁸. Bacteria were incubated in nutrient agar (NA-Merck) at 37 °C for 24 hours, and yeast was incubated for 48 hours. A few colonies were emulsified in 5 ml Nutrient Broth (NB-Merck), and adjusted spectrophotometrically to 0.1 absorbance at OD_{600} . The final inoculum was adjusted to 10^5 CFU/mL with 60 ml NB. From the resulting suspension, a 1.8 ml aliquot was transferred into first tube, added 1 ml into another tubes, then 200 μ L of metal complexes was transferred into the first tube. The samples were serially diluted for 1/2 (from 3000 to 5.9 μ g/ml). The samples were then incubated at 37 °C for 24 hours. The same study was repeated after narrowing the gap. The MIC was defined as the lowest concentration that completely prevented visible growth of the bacteria after overnight incubation at 37°C. The study was carried out in three replicates. Minimum Microbiocidal concentration (MMC) was determined only in the $Cu(pzdc)(phen)_2 \cdot 5.5H_2O$ complex. To determine minimum microbiocidal concentration (MMC) of $Cu(pzdc)(phen)_2 \cdot 5.5H_2O$ complex, 100 μ l from each tube which did not show growth in the MIC tests, and were incubated 37 °C for 24 h in nutrient agar medium. After incubation, the MBC value was determined as the lowest concentration that did not show growth in petri dishes.

Effect of $[Cu(pzdc)(phen)_2] \cdot 5.5H_2O$ complex on growth of bacteria

Each strain was grown in 5 ml NB, for 18

hours at 37 °C, in a shaking incubator at 175 rpm. After incubation, the cultures were adjusted spectrophotometrically to 0.1 absorbance at OD₆₀₀ and were used as starter cultures. The diluted overnight cultures (1:250 in NB) were incubated again (175 rpm, 37°C) until they reached an OD₆₀₀ 0.1, and defined concentrations of complexes (9 and 18 µg/ml) were added, incubated for 4 hours with the complex, and the turbidity of cultures growing in NB was determined at 600 nm at several time points by measuring the absorbance using an UV/Vis spectrophotometer (Thermo-Germany). The procedure was repeated three times. The average values were obtained.

Effect of [Cu(pzdc)(phen)₂].5.5H₂O on survival of microorganisms in lake water

The effect of [Cu(pzdc)(phen)₂].5.5H₂O complex on survival of bacteria and yeast in lake water was tested at different temperatures (37 °C and 10 °C) and at different concentrations (18, 9, 4.5 µg/ml). For this purpose, *E. coli*, *S. aureus*, *P. aeruginosa* and *C. albicans* were used. Bacteria were incubated in NB (Merck) at 37 °C for 24 hours, and yeast was incubated for 48 hours. The bacteria and yeast were added into 100 ml of filtered-autoclaved lake water (final concentration approximately 5×10⁵ CFU/ml for bacteria and 3×10⁴ CFU/ml for yeast), the complex was transferred into the lake water (final concentrations of 18, 9 and 4.5 µg/ml), and incubated at different temperatures (37 and 10 °C) for six hours. The samples were serially

diluted in Ringer's solution at several time points of incubation, including 240 minutes and plated onto NA. The colonies were counted after incubation at 37 °C for 24 h.

Data analyses

The JMP software was used for drawing graphics. Tests were carried out in independently three replicates.

RESULTS AND DISCUSSION

Resistant bacteria such as *S. aureus* and *P. aeruginosa* that are known to cause hospital infections have developed resistance against antibiotics¹⁹. New active substances are needed not only for human pathogens, but also for microorganisms that cause disease in plants and animals, and for disinfection of both water and environment. Therefore, the development of new antibiotics and new substances with antimicrobial activity is most urgent.

The antimicrobial activities of eight pyrazine-2,3-dicarboxylate complexes, which were previously synthesized and characterized^{14,15, 16, 17}, were investigated in this study, and the MICs of the complexes are presented in Table 3. [Cu(pzdc)(phen)₂].5.5H₂O clearly inhibited all test organisms at concentrations of 40 µg/ml, but the MIC for the clinical isolate *P. aeruginosa* was 100 µg/ml. The other seven complexes had values in the range of 400 - 3000 µg/ml against all the strains.

Table 1. Microorganisms used in this study

Microorganisms					
Wild type(Prokaryote)	Gram(+)	<i>S. aureus</i>	ATCC6538		
		<i>S. epidermidis</i>	ATCC12228		
		<i>B. cereus</i>	ATCC7064		
		<i>E. faecalis</i>	ATCC29212		
	Gram(-)	<i>P. aeruginosa</i>	ATCC27853		
		<i>E. coli</i>	ATCC25922		
		<i>S. typhimurium</i>	LT2		
		Clinical isolates(Prokaryote)	Gram(+)	MRSA-Methicillin resistant <i>S. aureus</i>	
				MRCNS- Methicillin resistant coagulase negative <i>Staphylococcus</i>	
Gram(-)	<i>P. aeruginosa</i>				
	<i>E. coli</i>				
	<i>M. morganii</i>				
Eukaryote			<i>A. baumannii</i>		
		<i>C. albicans</i>	ATCC 10231		

[Cu(pzdc)(phen)₂].5.5H₂O complex is active against both wild-type and clinically isolated bacteria. The effectiveness of this substance did not change importantly according to the cell wall structure of bacteria. Gram (-) bacteria, due to the presence of the outer membrane, are known to be more resistant compared to the Gram (+) ones²⁰. Some substances are not effective because they can not pass through the outer membrane of Gram (-) bacteria. However, they can be said to be more effective on the Gram-positive bacteria based on the MIC values of *S. aureus* (both wild-type and the clinical isolates, 15 µg/ml) and *B. cereus* (15 µg/ml). Furthermore, there are not any differences between prokaryotic and eukaryotic microorganisms in terms of the effectiveness of the [Cu(pzdc)(phen)₂].5.5H₂O complex.

[Cu(pzdc)(phen)₂].5.5H₂O was synthesized from pyrazine-2,3-dicarboxylic acid, 1,10-phenanthroline, and Cu(CH₃COO)₂.H₂O. If we look at the results of Table 3 individually, the MIC values of pyrazine-2,3-dicarboxylic acid for six microorganisms were 312.5 µg/ml and for eight microorganisms, the value was 625 µg/ml. In addition, the MIC values of 1,10-phenanthroline monohydrate were in the range 15-312.5 µg/ml and CuCl₂.2H₂O were in the range 312.5-625 µg/ml for all tested bacteria and yeast. The complex is also highly active compared to the parent compounds (from 9 and 10) shown by the 8- to 64-fold lower MIC value (Table 3). [Cu(pzdc)(phen)₂].5.5H₂O was more potent than pyrazine-2,3-dicarboxylic acid¹⁰ in all cases and nearly equipotent with the 9 against *E. coli*, *S. typhimurium*, *B. cereus* and some clinical isolates (*E. coli*, *A. baumannii*, *M. organii*, and *C. albicans*). The synthesized complex was found to have lower MIC value according to the ligands and the main matter. Our results clearly showed that [Cu(pzdc)(phen)₂].5.5H₂O possesses broad antibacterial and anticandidal effects against the microorganisms used in the study. After this study, the MIC values of [Cu(pzdc)(phen)₂].5.5H₂O complex were determined that it did not kill microorganisms both using plate count and microscope. Moreover, the minimum microbiocidal concentration (MMC) of [Cu(pzdc)(phen)₂].5.5H₂O was investigated, MMC values are compared in Table 3. There is a big difference between the MIC (15-40 µg/ml) and MMC (30-200 µg/ml) values. Therefore, the mechanism of action of this complex

is thought to be associated with both cell wall integrity and replication mechanisms.

The bacteriocidal potential of [Cu(pzdc)(phen)₂].5.5H₂O was assessed in time-kill assays with one Gram-positive organisms (*S. aureus*) and two Gram-negative bacteria (*E. coli* and *P. aeruginosa*). According to Pankey and Sabath²¹, a compound is defined as bacteriocidal if it effects a 3-log or greater decrease in bacterial cell density after 24 hours of incubation. According to the results of the MIC, the effect of the [Cu(pzdc)(phen)₂].5.5H₂O complex, which was determined to be highly effective in NB, its effect on survival of microorganisms in lake water was examined. The effect of temperature on the activity of the complex, the effect of different concentrations of the complex on its antimicrobial activity, and its effectiveness against the microorganisms in lake water as a natural environment were investigated. The data obtained from these experiments are presented in the Figures 2-6. The lake-water experiments showed that temperature is very important for the effectiveness of complex. While *E. coli*, *S. aureus* and *P. aeruginosa* were completely inhibited at a concentration of 4.5 µg/ml at 37 °C in a period of 240 minutes, these bacteria were not inhibited even at a concentration of 18 µg/ml at 10 °C in the same time interval (Fig. 3-5). Another interesting observation encountered in this study was the effect of the complex on the survival of *C. albicans*

Table 2. Metal complexes used in this study

	Complexes and Ligant	Ref.
1	[Cu ₂ (μ-pzdca) ₂ (pen) ₂].2H ₂ O	16
2	{[Cu(μ-pzdca)(bipy)]·H ₂ O} _n	16
3	[Cu(pzdca)(H ₂ O)(dmpen) ₂]	15
4	{[Cu(μ-pzdca)(tmen)]·H ₂ O} _n	16
5	[Cu(pzdca)(H ₂ O)(en) ₂].H ₂ O	16
6	[Cu(μ3-pzdca)(mea)] _n	17
7	[Cu ₂ (pzdca) ₂ (H ₂ O) ₂ (dmen) ₂].6H ₂ O	14
8	[Cu(pzdca)(phen) ₂].5.5H ₂ O	14
9	1,10 phenanthroline monohydrate	14
10	Pyrazine-2,3-dicarboxylic acid (ligand)	14

[Abbreviations: pzdca=Pyrazine-2,3-dicarboxylic acid, pen=1,3-propanediamine, bipy=2,2'-bipyridine, dmpen=2, 2-dimethylpropane-1,3-diamine, tmen=N,N,N',N'-tetramethylethylenediamine, en=ethylenediamine, dmen=N,N'-dimethylethylenediamine, phen=1,10-phenanthroline, mea= monoethanolamine]

Table 3. Minimal inhibition concentrations (MIC) and Minimum microbiocidal concentration values (MMC of 8) of complexes and ligand for wild type and clinical isolate ($\mu\text{g/ml}$)

Microorganisms		1	2	3	4	5	6	7	8	9	10	11	
		(MIC)	(MIC)	(MIC)	(MIC)	(MIC)	(MIC)	(MIC)	(MIC)	(MMC)	(MIC)	(MIC)	
Wildtype	Gram(-)	<i>E. coli</i>	800	600	1400	1500	2500	1400	<3000	40	<100	30	625
		<i>P. aeruginosa</i>	1500	1250	1400	1500	2750	<3000	<3000	20	100	312.5	625
		<i>S. typhimurium</i>	900	900	2250	1750	<3000	2500	<3000	30	<100	30	625
	Gram(+)	<i>E. faecalis</i>	1500	750	1500	1000	<3000	<3000	<3000	20	60	312.5	312.5
		<i>B. cereus</i>	1000	750	2500	1500	2750	2500	<3000	15	30	20	312.5
Clinicalisolates	Gram(+)	<i>S. epidermidis</i>	1000	750	2500	1500	2000	2750	<3000	30	40	156	625
		<i>S. aureus</i>	800	750	1000	1500	1250	1700	<3000	15	30	250	625
		<i>MRSA</i>	1250	750	1500	1750	<3000	1700	<3000	15	40	75	312.5
		<i>MRCNS</i>	600	500	750	1000	1000	2500	<3000	20	60	40	625
		<i>E. coli</i>	800	1500	1750	1750	<3000	1400	<3000	35	70	30	625
Eucaryote	Gram(-)	<i>P. aeruginosa</i>	1500	1500	<3000	<3000	<3000	<3000	<3000	100	<200	312.5	625
		<i>A. baumannii</i>	700	1250	2750	3000	1500	2500	<3000	35	60	40	625
		<i>M. morganii</i>	1500	400	1750	<3000	2750	<3000	<3000	30	<100	40	625
		<i>C. albicans</i>	<3000	<3000	<3000	<3000	<3000	<3000	<3000	25	60	30	625
		2- { [Cu (<-pzdea)(bipy)] · H ₂ O } _n , 5- [Cu (pzdea)(H ₂ O)(en) ₂] · H ₂ O, 8- [Cu (pzdea)(phen) ₂] · 5.5H ₂ O, 11- CuCl ₂ · 2H ₂ O											
		3-[Cu(pzdea)(H ₂ O)(dmpen)2], 6-[Cu (<-pzdea)(mea)] _n , 9-1,10-phenanthroline monohydrate (phen),											




Fig. 1. Structure of the $[\text{Cu}(\text{pzdc})(\text{phen})_2] \cdot 5.5\text{H}_2\text{O}$ complex²⁷




Fig. 2. Inhibition of the growth of microorganisms with the $[\text{Cu}(\text{pzdc})(\text{phen})_2] \cdot 5.5\text{H}_2\text{O}$ complex. The complex was added to early logarithmic phase cultures at 0. hour (approximately 0.1 at 600 nm) (A- *E. coli*, B- *S. aureus*, C- *C. albicans*, D- *P. aeruginosa*)




Fig. 3. Effect of the $[\text{Cu}(\text{pzdc})(\text{phen})_2] \cdot 5.5\text{H}_2\text{O}$ complex on survival of *E. coli* in lake water

(Fig. 6). *C. albicans* was not inhibited even at 18 µg/ml, at 37 °C, in period of 240 minutes. None of the studied conditions showed any remarkable effect on *C. albicans* in lake water (Fig. 6). This results is quite interesting.

A concentration of 4.5 µg/ml of this complex in lake water showed a 3 log. decline within 45 minutes for *E. coli* and *S. aureus*, and in 180 minutes for *P. aeruginosa*. These results indicate that while 18 µg/ml of this complex had a bacteriostatic effect in NB, the same concentration had a bactericidal effect on microorganisms in lake

water. When the activity of $[\text{Cu}(\text{pzdc})(\text{phen})_2] \cdot 5.5\text{H}_2\text{O}$ complex was compared at two temperature, 37 and 10 °C, the complex was found to be very effective at 37 °C than at 10 °C. The reason for this effect is thought to be a the high levels of metabolic activity reached by microorganisms in their logarithmic growth phase, in addition to the temperature 37 °C being the optimum temperature required for the growth of microorganisms. The effectiveness of the complex is thought decreased at 10 °C due to the slower metabolism of the microorganisms.




Fig. 4. Effect of the $[\text{Cu}(\text{pzdc})(\text{phen})_2] \cdot 5.5\text{H}_2\text{O}$ complex on survival of *S. aureus* in lake water




Fig. 5. Effect of the $[\text{Cu}(\text{pzdc})(\text{phen})_2] \cdot 5.5\text{H}_2\text{O}$ complex on survival of *P. aeruginosa* in lake water




Fig. 6. Effect of the $[\text{Cu}(\text{pzdc})(\text{phen})_2] \cdot 5.5\text{H}_2\text{O}$ complex on survival of *C. albicans* in lake water

When the $[\text{Cu}(\text{pzdc})(\text{phen})_2] \cdot 5.5\text{H}_2\text{O}$ complex is dissolved in water, it is composed of a mononuclear structure consisting of a Cu(II) ion, two 1,10-phenanthroline ligands and one pyrazine-2,3-dicarboxylate dianion. When the structure-activity relationship of the complex was examined in isolation, the MIC values of 1-10-phenanthroline is higher, in general, than metal complex. Yesilel *et al.*²² reported that the MIC value of $[\text{Co}(\text{pzdc})(\text{phen})_2]_2 \cdot 11\text{H}_2\text{O}$ was 2000-2250 $\mu\text{g}/\text{ml}$. $[\text{Co}(\text{pzdc})(\text{phen})_2]_2 \cdot 11\text{H}_2\text{O}$ and $[\text{Cu}(\text{pzdc})(\text{phen})_2] \cdot 5.5\text{H}_2\text{O}$ have an identical molecular structure. Therefore, the difference between the two molecules is only that the current compound under study has Cu(II) instead of Co(II) as the central metal moiety. However, the effectiveness of the molecule cannot be said to be derived completely either due to Cu(II) or phenanthroline (Table 3). The structure of $[\text{Cu}(\text{pzdc})(\text{phen})_2] \cdot 5.5\text{H}_2\text{O}$ as a whole can be said to have a strong effect on microorganisms. In lake water studies, with an MIC value of 20 $\mu\text{g}/\text{ml}$ against *C. albicans*, it was found to be virtually ineffective because *C. albicans* is in the logarithmic growth phase in NB and in the stationary phase in lake water.

2,3-diethyl pyrazine dicarboxylic acid which was produced from pyrazine dicarboxylic acid was found to prevent corrosion in steel²³. The pyrazine ring is known to be a part of many polycyclic compounds of biological and industrial importance. In addition, pyrazine derivatives have been reported to possess antituberculous, antifungal, and cytotoxic activities²⁴⁻²⁶. Especially, pyrazine-2,3-dicarboxylic acid derivatives were shown to be effective on *Mycobacterium tuberculosis* which causes the dreaded tuberculosis disease in the third-world countries and is beginning to be a problem again currently²⁷⁻²⁹. Pyrazine derivatives were seen as suitable candidates for the antimicrobial substances³⁰. In addition, in some studies of the antimicrobial activity of various compounds, the hydrazinium salts of 2-pyrazinecarboxylate, pyrazine-2,3-dicarboxylate, pyrazine-3,5-dicarboxylate and imidazole-4,5-dicarboxylate have been found to be more effective than standard antibiotics against *E. coli*, *Salmonella typhi* and *Vibrio cholerae*^{31,32}. In addition, pyrazine-2-carboxylic acid derivatives inhibited the growth of *Leuconostoc mesenteroides*³³.

According to the obtained data of the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) cytotoxicity test, $[\text{Cu}(\text{pzdc})(\text{phen})_2] \cdot 5.5\text{H}_2\text{O}$ was found to have a IC_{50} value of 5 $\mu\text{g}/\text{ml}$ against NS20Y neuroblastoma cells (detailed data did not given).

Thus, the $[\text{Cu}(\text{pzdc})(\text{phen})_2] \cdot 5.5\text{H}_2\text{O}$ complex is determined to be very effective even at low concentrations with 15-40 $\mu\text{g}/\text{ml}$ being the MIC value against microorganisms. This complex can be applied as a disinfectant because of its very low MIC value, is very easily soluble in water, and is odorless and colorless at the effective concentration. This novel metal complex which is potent against antibiotic-resistant pathogens, should be further investigated. Combining the information obtained in this study with pharmacokinetic and pharmacodynamic data is necessary to provide an in vivo efficacy prediction that is more meaningful.

REFERENCES

1. Dennesen, P.J., Bonten, M.J., Weinstein, R.A. Multiresistant bacteria as a hospital epidemic problem, *Ann. Med.*, 1998; **30**: 176-85
2. Hancock, R.E., Speert, D.P. Antibiotic resistance in *Pseudomonas aeruginosa*: mechanisms and impact on treatment. *Drug Resist Update*, 2000; **3**: 247-55.
3. Kaye, K.S., Engemann, J.J., Fraimow, H.S., Abrutyn, E. Pathogens resistant to antimicrobial agents: epidemiology, molecular mechanisms, and clinical management. *Infect Dis. Clin. North. Am.*, 2004; **18**: 467-11.
4. Prystowsky, J., Siddiqui, F., Chosay, J., Shinabarger, D.L., *et al.*, Resistance to linezolid: characterization of mutations in rRNA and comparison of their occurrences in vancomycin-resistant enterococci. *Antimicrob. Agents Chemother.*, 2001; **45**: 2154-6.
5. Roberts, M.C. Antibiotic resistance in oral/respiratory bacteria, *Crit. Rev. Oral. Biol. Med.*, 1998; **9**: 522-40.
6. Wang, E.E., Kellner, J.D., Arnold, S. Antibiotic-resistant *Streptococcus pneumoniae*. Implications for medical practice, *Can. Fam. Physician*, 1998; **44**: 1881-8.
7. Ar, C. Protection from Hospital Infections in immunocompromised patients, Cerrahpasa Medical Faculty, Medical Education Activities, Hospital Activities: *Symposium Series*, 2008; **60**: 239-53.

8. Baran, E.J., Yilmaz, V.T. Metal complexes of saccharin, *Coor. Chem. Reviews*, 2006; **250**: 1980-99.
9. Yesilel, O.Z., Kastan, G., Darcan, C., Ilker, I., Pasaoglu, H., Buyukgungor, O. Syntheses, thermal analyses, crystal structures and antimicrobial properties of silver(I)-saccharinate complexes with diverse diamine ligands, *Inorganica Chimica Acta*, 2010; **363**: 1849-58.
10. Olar, R., Badea, M., Marinescu, D., Chifiriuc, C.M., et al., Prospects for new antimicrobials based on N,N-dimethylbiguanide complexes as effective agents on both planktonic and adhered microbial strains. *Eur. J. Med. Chem.*, 2010; **45**: 2868-75.
11. Rai, M., Yadav, A., Gade, A. Silver nanoparticles as a new generation of antimicrobials, *Biotechnology Adv*, 2009; **27**: 76-3.
12. Landsdown, A.B. Silver. I: Its antibacterial properties and mechanism of action *J. Wound Care*, 2002; **11**: 125-30.
13. Hussain, S., Ferguson, C. Best evidence topic report. Silver sulphadiazine cream in burns., *Emerg Med. J.*, 2006; **23**: 929-32.
14. Yesilel, O.Z., Mutlu, A., Buyukgungor, O. A new coordination mode of pyrazine-2,3-dicarboxylic acid and its first monodentate complexes: Syntheses, spectral, thermal and structural characterization of [Cu(pzdc)(H₂O)(en)]·H₂O and [Cu(pzdc)(H₂O)(dmpen)]₂, *Polyhedron*, 2008; **27**: 2471-77.
15. Yesilel, O.Z., Mutlu, A., Büyükgüngör, O. Syntheses, spectral, thermal and structural characterization of copper(II) complexes of 2,3-pyrazinedicarboxylate with N, N 2 -dimethylethylenediamine and 1,10-phenanthroline, *J. Coord. Chem.*, 2008; **61**: 3267-75.
16. Yesilel, O.Z., Mutlu, A., Büyükgüngör, O. Novel dinuclear and polynuclear copper(II)-pyrazine-2,3-dicarboxylate supramolecular complexes with 1,3-propanediamine, N,N,N₂,N₂ -tetramethylethylenediamine and 2,22-bipyridine *Polyhedron*, 2009; **28**: 437-44.
17. Yesilel, O.Z., Mutlu, A., Günay, G., Caner, N., et al, Copper(II)-pyrazine-2,3-dicarboxylate coordination polymer with monoethanolamine, [Cu(μ₃-pzdc)(mea)]_n *J. Chem. Crystallogr.*, 2011; **42**: 519-23.
18. NCCLS guideline. Clinical and Laboratory Standards Institute 8th ed. Document M7-A8. Wayne, PA: CLSI 2009.
19. Novick, R.P., Schlievert, P., Ruzin, A. Pathogenicity and resistance islands of staphylococci, *Microbes Infect.*, 2001; **3**: 585-95.
20. Nikaido, H. Outer membrane barrier as a mechanism of antimicrobial resistance. *Antimicrob. Agents Chemother.*, 1989; **33**: 1831-6.
21. Pankey, G.A., Sabath, L.D. Clinical relevance of bacteriostatic versus bactericidal mechanisms of action in the treatment of gram-positive bacterial infections, *Clin. Infect. Dis.*, 2004; **38**: 864-70.
22. Yesilel, O.Z., Mutlu, A., Darcan, C., Buyukgungor, O. Syntheses, structural characterization and antimicrobial activities of novel cobalt-pyrazine-2,3-dicarboxylate complexes with N-donor ligands, *J. Mol. Struct.*, 2010; **964**: 39-46;
23. Bouklah, M., Attayibat, A., Kertit, S., Ramdani, A., et al., A pyrazine derivative as corrosion inhibitor for steel in sulphuric acid solution, *Applied Surface Science*, 2005; **242**: 399-406.
24. Adams, T.B., Doull, J., Feron, V.J., Goodman, J.I., Marnett, L.J., Munro, I.C., et al., The FEMA GRAS assessment of pyrazine derivatives used as flavor ingredients. Flavor and Extract Manufacturers Association, *Food Chem. Toxicol.* 2002; **40**: 429-51.
25. Dolezal, M., Jampilek, J., Osicka, Z. Substituted 5-arylpyrazine-2-carboxylic acid derivatives: synthesis and biological activity, *Farmaco*, 2003; **58**: 1105-11.
26. Seitz E.L., Suling, J.W., Reynolds, C.R. Synthesis and Antimycobacterial Activity of Pyrazine and Quinoxaline Derivatives, *J. Med. Chem.*, 2002; **45**: 5604-6.
27. Abdel-Aziz, M., Abdel-Rahman, H.M. Synthesis and anti-mycobacterial evaluation of some pyrazine-2-carboxylic acid hydrazide derivatives, *Eur. J. Med. Chem.*, 2010; **4**: 3384-8.
28. Dolezal, M., Cmedlova, P., Palek, L. Synthesis and antimycobacterial evaluation of substituted pyrazinecarboxamides, *Eur. J. Med. Chem.*, 2008; **43**: 1105-13.
29. Zimhony, O., Vilcheze, C., Arai, M. Pyrazinoic Acid and Its n-Propyl Ester Inhibit Fatty Acid Synthase Type I in Replicating Tubercle Bacilli, *Antimicrob. Agents Chemother.*, 2007; **51**: 752-4.
30. Buczkowska, M., Bodtke, A., Lindequist, U., Gdaniec, M., Bednarski, P.J. Cytotoxic and antimicrobial Activities of Cu(II), Co(II), Pt(II) and Zn(II) complexes with N,O-Chelating heterocyclic carboxylates, *Arch. Pharm. Chem. Life Sci.*, 2011; **344**: 605-16.
31. Premkumar, T., Govindarajan, S. Antimicrobial study of pyrazine, pyrazole and imidazole

- carboxylic acids and their hydrazinium salts, *World J. Microbiol. Biotech.*, 2005; **21**: 479-80.
32. Premkumar, T., Govindarajan, S. Antimicrobial study on trivalent lighter rare-earth complexes of 2-pyrazinecarboxylate with hydrazinium cation, *World J. Microbiol. Biotech.*, 2006; **22**: 1105-8.
33. Abd El-Wahab, A.H.F., Bedair, A.H., Eid, F.A., El-Haddad, A.F., Adawy El-Deeb, A.M., El-Sherbiny, G.M. Pyrazine-2-substituted carboxamide derivatives: synthesis, antimicrobial and *Leuconostoc mesenteroides* growth inhibition activity, *Journal of the Serbian Chemical Society*, 2006; **71**: 471-81.