Beautlin et al | Article 8752 J Pure Appl Microbiol. 2023;17(4):2157-2164. doi: 10.22207/JPAM.17.4.12 Received: 05 June 2023 | Accepted: 04 September 2023 Published Online: 23 October 2023

RESEARCH ARTICLE



Comparing hs-CRP and MPO Biomarkers of Subclinical Atherosclerosis in Metabolic Syndrome Patients

Abishkauf Jenish Beautlin^{1*}, Ashok Govindaraj¹, Durga Devi¹, Gurubharath Ilangovan² and Divya Lakshmi¹

¹Department of Cardiology, Chettinad Academy of Research and Education, Kelambakkam, Tamil Nadu, India. ²Department of Radiology, Chettinad Academy of Research and Education, Kelambakkam, Tamil Nadu, India.

Abstract

Cardiovascular disorders have been linked to myeloperoxidase, an enzyme generated from leukocytes. MPO creates an antibacterial system and has been many correlations between MPO, hs-CRP and cardiovascular disease. The present study has aimed to assess MPO, hs-CRP and subclinical atherosclerosis in metabolic syndrome patients by using FMD and CIMT parameters. 75 metabolic syndrome affected patients had defined as per the criteria of IDF (International Diabetes Federation) which had been in this study. In addition, ultrasound doppler had been employed for determining the carotid intima medial thickness for both right and left (left and right CIMT) and brachial artery FMD. By employing the Immunoassay MPO and hs-CRP ultrasensitive ELISA kit, the concentration of MPO and hs-CRP is measured. A positive link between MPO with CIMT right and left (r value of right CIMT=0.723, p <0.05 and left CIMT r=0.712, p<0.01), hs-CRP with CIMT right and left (r value of right CIMT=0.613, p<0.05and left CIMT r=0.64, p<0.01)and a significant inverse correlation between MPO with FMD (r= -0.319,p<0.05), hs-CRP with FMD (r= -0.304,p<0.01) in metabolic syndrome patients. Serum MPO and hs-CRP is positively linked with subclinical atherosclerosis in metabolic syndrome patients. According to this finding, understanding the mechanism behind the risk factors for cardiovascular disease can be aided by the development of new molecular markers for metabolic syndrome and subclinical atherosclerosis.

Keywords: Flow Mediated Vasodilatation, Carotid Intima Medial Thickness, Metabolic Syndrome, hs-CRP, Myeloperoxidase

*Correspondence: rajjenish1996@gmail.com

Citation: Beautlin AJ, Govindaraj A, Devi D, Ilangovan G, Lakshmi D. Comparing hs-CRP and MPO Biomarkers of Subclinical Atherosclerosis in Metabolic Syndrome Patients. *J Pure Appl Microbiol.* 2023;17(4):2157-2164. doi: 10.22207/JPAM.17.4.12

© The Author(s) 2023. **Open Access**. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License which permits unrestricted use, sharing, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

Journal of Pure and Applied Microbiology

INTRODUCTION

Syndrome X or the Metabolic syndrome (MS) from the clinical standpoint represents the combination of various metabolic abnormalities that contribute increased cardiovascular disease.¹ In CVD related complication, atherosclerosis is regarded as a systematic, chronic and progressive condition observed with one of the prolonged asymptomatic phases.² Due to the macrophages activation and cytokines release by expanded adipocytes, pro-inflammatory state serves a pivotal role in the syndrome pathogenesis. Estimation of C-reactive protein level in serum can document this pro-inflammatory state.^{3,4} From the point of fact, the criteria were initially proposed by WHO to define Metabolic Syndrome, which has been evolved over the course of years. Currently, Metabolic Syndrome criteria remain the basis of IDF (International Diabetes Federation) criteria, multiple clinical &laboratory parameters constellation. Since there is a dire need for predicting CVD related complications,⁵⁻¹⁰ the recent interest in the investigation on numerous biomarkers and their influence in the prediction of cardiovascular events could greatly facilitate in devising a suitable model that compares clinical variables and patient's demographics.¹¹⁻¹³ High sensitivity CRP levels need to be evaluated in metabolic syndrome.14-18 The level of HDL and obesity had been regarded out of all metabolic syndrome features. High sensitivity CRP levels had been thoroughly analysed in obesity patients.^{19,20}

Gradually, atherosclerosis has developed as one of the subclinical conditions over the course of life and they eventually become apparent as peripheral arterial disease clinically.²¹⁻²³ The early stage atherosclerosis is referred as pre- or sub-clinical atherosclerosis, when it is within the vascular walls that is, something starts to get changed, but the cardiovascular disease isn't evident clinically (Figure). Interest over the past decade has been gained by detecting the upcoming disease before the clinical manifestations at this stage.^{24,25} As a result, early detection and intervention of the progression of atherosclerosis is critical for minimising the burden of CVDs and stroke in the general population.²⁶⁻³⁰ Carotid intimamedia thickness (CIMT) ultrasonography measures give a good indication of early atherosclerosis that is independent of established risk factors, and the CIMT is thought to be beneficial in predicting cardiovascular events including ischemic strokes.^{31,32} Chronic inflammation is thought to play a role in the development of cardiovascular illnesses (CVD), such as coronary heart disease and stroke, as well as metabolic disorders.³⁴⁻³⁶ Three subclinical atherosclerosis measures remain coronary artery calcification, ankle-branchial index & carotid intima-media thickness which can possibly be quantified as well as non-invasively detected.

MATERIALS AND METHODS

The investigation was done in Chettinad Hospital and Research Institute, Kelambakkam. The period of study is extended between 2021 and 2022. All patients gave their informed consent before participating. The study population comprised of patients presented with metabolic syndrome were carefully selected on the basis of IDF (International Diabetes Federation) criteria. Based on the inclusion and exclusion criteria, every patient had been selected. High-sensitivity CRP and MPO had been estimated and analysed in all those patients. Both gender patients had been included in the study. About 75 patients had been included in this study. Predominantly, the sample had been collected from the patients who attended the master health check-up.

This investigation is a cross sectional study. Randomly, the study had been chosen from the master health examination patients. For metabolic syndrome, many quarters have approved and accepted the IDF guidelines. It has been the one that extensively employed in the clinical metabolic syndrome trials. 75 clinical samples are collected from metabolic syndrome patients.

The quantitative measurement was carried out using the enzyme-linked immunosorbent assay of MPO and hs-CRP in serum samples. Further, readings had been measured. As per the kit instructions, the procedure of ELISA had been performed.

Subclinical atherosclerosis parameters:

Experience radiologist took CIMT and FMD in the department of radiology. Metabolic syndrome – there is the presence of more than or equal to three in the IDF criteria. Raised triglyceride:>150mg/dl

Reduced HDL cholesterol:<40mg/dl in males and <50 mg/dl in females

Raised Blood Pressure: Systolic BP>130 or diastolic BP >85mm Hg

Raised fasting plasma glucose: FBG>100 mg/dL Central obesity (Waist circumference) :(>80cm female) and (>90cm male)

Inclusion criteria

The patient must meet the criteria for metabolic syndrome as outlined in the IDF recommendations previously mentioned.

Exclusion criteria

- 1. Previous record of coronary artery disease
- 2. Previous record of cerebrovascular accidents.
- Patients with inflammation, neoplasm, pregnancy smokers, autoimmune, hepatic and renal dysfunction and terminal illness were excluded from the study.

FMD analysis

Assessment of brachial FMD was conducted among the study population in their fasting state. All the participants were instructed to lie down in a supine position for at least 10 minutes before the start of the test. A blood pressure cuff is placed around the participant's forearm, and an ultrasound probe is placed on the arm to visualize the brachial artery. The blood pressure cuff is inflated to a pressure of 200 mmHg to induce ischemia in the forearm. The cuff is then rapidly deflated, which leads to a reactive hyperaemic response that causes an increase in blood flow to the arm and a corresponding increase in brachial artery diameter. 70 to 90 seconds after the cuff deflates, the diameter of the brachial artery is measured. The FMD is calculated using the formula as follows: (maximum diameter – baseline diameter)/baseline diameter ' 100%.

If the FMD was less than 10%, endothelial dysfunction was declared to be the cause.

CIMT

CIMT (carotid intima-media thickness) is a safe procedure for assessing subclinical form of atherosclerosis. The methodology for CIMT measurement typically involves the following steps:

The participant is asked to lie down in a supine position, and the neck is slightly extended. An ultrasound probe with a high-frequency



Figure. Progression of atherosclerosis

transducer (7-10 MHz) is placed on the skin over the carotid artery. The ultrasound probe is used to see the intimate-media complex of the carotid artery. The intima-media complex is the two innermost layers of the artery wall, which consist of the intima (the innermost layer) and the media (the middle layer). The CIMT is measured as the distance between the lumen-intima interface and the media-adventitia interface on both the left and right sides of the carotid artery. Measurements are taken at several points along the length of the carotid artery, typically at the common carotid, carotid bulb, and internal carotid segments. The mean CIMT is calculated as the average of the measurements taken at each segment of the carotid artery.

Statistical analysis

SPSS IBM software version 26 software was used for data analysis. The mean and standard

Tak	ole :	1.	Baseline	characteristics	of	the	patients
-----	-------	----	----------	-----------------	----	-----	----------

Parameters	Mean ± Std. Deviation (Range)
Age (years)	58.08±10.247 (36-85)
FBS in plasma	145.05±50.720
(Normal 70–110 mg/dL)	
TGL (Normal < 150 mg/ dL)	194.79±115.074
HDL (mg/ dL)	37.13 ±7.780
Waist Circumference	77.72 ±10.147
(inches)	
SBP (mm Hg)	120.24±10.917
DBP (mm Hg)	81.40±10.351
CIMT-Right (mm)	1.0467±0.25697
CIMT-Left (mm)	1.0160±0.31151
FMD(%)	9.5240±2.514
hs-CRP (mg/ L)	2.5467±1.328
MPO (AU/mL)	26.66 ±6.96

Table 2. Correlation of MPO with CIMT: Considerably enhanced the diameter of the right and left CIMT as compared to the MPO with CIMT

			_
MPO	CIN	ИТ	
	RIGHT	LEFT	
26.66 ± 6.96	1.04±0.25**	1.01±0.31**	

Values expressed as mean standard deviation with pearson correlation significant levels of **p value <0.01 (r value right CIMT=0.723 and left CIMT r=0.712)

deviation (SD) were used to represent continuous variables. Correlation between the biomarkers were calculated by using Pearson's correlation. The independent t-test was used to compare two independent populations. P-value <0.05 was taken as statistical significance.

RESULTS

Study has 75 patients of which 47 were male (62.7%) and 28 were women (37.3%) with a mean age of 58.0 ± 10.2 years. Table 1 shows the Abnormal FBS ranges (N=69; 92%) with a mean average of 145.05 ± 50.720 . In HDL (N=62;82.7%) patients were presented with abnormal with a mean average of 37.13 ± 7.780 . Abnormal ranges in TGL (N=50; 66.6%) with a mean average of 194.79 \pm 115.074. Abnormal ranges in waist

 Table 3. Correlation of hs-CRP with CIMT: Compared to the HS-CRP with CIMT, the right and left CIMT's diameters are significantly increased

Hs-CRP	CIMT		
	RIGHT	LEFT	
2.54±1.32	1.04±0.25**	1.01±0.31**	:

Values expressed as mean standard deviation with pearson correlation significant levels of *p value <0.01 (r value of right CIMT=0.613 and left CIMT r=0.641)

Table 4. Correlation of MPO with FMD using PearsonCorrelation: Inverse correlation exists between theMPO and FMD

MPO	FMD	
26.6±6.96	9.52±2.51*	

Values expressed as mean standard deviation with pearson correlation significant values of *p value<0.05* (r= -0.319)

 Table 5. Correlation of hs-CRP with FMD using Pearson

 Correlation Inverse correlation exists between the hs

 CRP and FMD

Hs-CRP	FMD	
2.54±1.32	9.52±2.51*	

Values expressed as mean standard deviation with pearson correlation inversely correlated (r= -0.304)

Journal of Pure and Applied Microbiology

www.microbiologyjournal.org

circumference (N=23; 30.6%). Blood pressure in abnormal ranges (N=34;45.3%).

DISCUSSION

A measure of subclinical atherosclerosis called CIMT is reliable predictor of cardiovascular and cerebral vascular events. Carotid ultrasonography has been shown in studies to be a sensitive technique that produces trustworthy results.³⁷ Furthermore, it has been demonstrated that the onset of clinically discernible atherosclerotic plaques in the coronary arteries precedes, which is recognised as first event in atherogenesis. Nitric oxide (NO), in particular, has a lower bioavailability due to endothelial dysfunction, although the bioavailability of contracting substances originating from the endothelium is increased. Many studies evaluated the endothelial function as a predictor of CVD.³⁸

A high concentration of MPO is thought to be a predictor of future cardiovascular disease in healthy individuals since inflammation is activated many years before the beginning of cardiovascular disease. Even after adjusting for conventional risk variables and hypertension, increased the levels of MPO and hs-CRP were independently identified a predictor of endothelial dysfunction and cardiovascular disease.³⁹

According to recent research, hs-crp and serum MPO levels were predictive of subclinical atherosclerosis. As a result, increasing CIMT in MS patients was positively linked with higher levels of MPO and hs-CRP (Table 2 and 3). According to Toprak and colleagues, who discovered that any rise in hs-CRP was connected to the advancement of CIMT regardless of other cardiovascular risk factors, there is a positive link between CIMT and hs-CRP.⁴⁰ In the INVADE trial, which examined the relationship between MPO and CIMT development in aged patients, the two parameters were found to be independently correlated.⁴¹ Andreeaiana et al. found that greater serum MPO levels, coupled with other biomarkers (C-reactive protein, plasminogen activator inhibitor-1, leptin, apolipoprotein-B, apolipoprotein-A1), predicted subclinical atherosclerosis. Thus, higher MPO levels were associated with increased CIMT in MS patients.

According to Qiaoxiayang et al., Metabolic syndrome components such as increased BP and abdominal obesity were positively associated with CIMT, but higher TG levels were adversely associated with CIMT.^{42,43} Previous cross-sectional investigations have shown that those with MetS have higher CIMTs.⁴⁴ Obesity is associated with a chronic inflammatory process characterised by an increase in different proinflammatory biomarkers, which may mediate all stages of atherogenesis, including the beginning, acceleration, and development of atherosclerotic lesions.⁴⁵ According to the Brazilian Longitudinal Study of Adult Health, for each 1-cm rise in WC, CIMT increased by 0.08 mm (= 0.08; 95% CI, 0.04-0.11; P 0.0001).46

hsCRP levels were also considerably higher in a subset of patients with CVD, hypertension, diabetes, and visceral obesity compared to a subset of people without such diseases.⁴⁷

Our study demonstrates a substantial relationship between serum MPO and hs-CRP and alterations in FMD in brachial arteries in Table 4 and 5. The consumption of nitric oxide, which may indicate a problem with endothelial function, is a significant effect of increased MPO activity. The endothelial dysfunction, was extensively evaluated in numerous research as a predictor of future cardiovascular events.^{48,49} A loss of the unadjusted relationship between C-reactive protein and endothelial dysfunction after risk factor and cardiovascular disease adjustment. Due to these complicating factors, the previously established association between C-reactive protein and endothelial dysfunction may still hold true.⁵⁰⁻⁵²

Our findings revealed favourable association between the biomarker ranges and subclinical atherosclerosis parameters in MS patients. This further suggest that the biomarkers could serve as a prognostic tool in detecting the complications and risks associated with CVDs that may contribute in the development of subclinical atherosclerosis. The study ultimately led us to the conclusion that the effects of hs-CRP and MPO could determine subclinical atherosclerosis condition. This showed that a simple blood test can be performed by any standard clinical laboratory and the versatility and possible clinical utility might be offered by MPO and hs-CRP testing in

Journal of Pure and Applied Microbiology

the disease advancement's serial monitoring. It is further suggested that subclinical atherosclerosis and new biological markers for MS have been helpful in order to understanding the mechanisms and the role.

CONCLUSION

Findings shows the serum biomarkers viz, MPO and hs-CRP were significantly correlated with subclinical atherosclerosis in patients with MS. It was indicated from the available data that hs-CRP selective determination has been beneficial in every individual with intermediate cardiovascular risks for optimizing the clinical management as well as risk stratification, in spite of limitations to the use of routine clinical practice, in particular, interindividual variability. In addition, on the basis of sound experimental evidences from the investigation linking with serum biomarkers with development of atherosclerosis represent a stronger association from the brachial FMD and CIMT measures observed in the current study, and earlier demonstration and promising as a biomarker target of subclinical atherosclerosis from the significant correlation in MPO and hs-CRP with subclinical atherosclerosis among metabolic syndrome affected patients. Cardiovascular risk stratification in metabolic syndrome affected patients with metabolic syndrome may be improved by the combination of ultrasonography alongside biomarker in association with vulnerable plaque pathophysiology namely MPO and hs-CRP. It is concluded that there is a strong relationship with regard to the elevated ranges in the biomarkers MPO and hs-CRP and the likely development in subclinical atherosclerosis condition among metabolic syndrome patients.

ACKNOWLEDGMENTS

None.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

AUTHORS' CONTRIBUTION

All authors listed have made a substantial, direct and intellectual contribution to the work, and approved it for publication.

FUNDING

None.

DATA AVAILABILITY

All datasets generated or analyzed during this study are included in the manuscript.

ETHICS STATEMENT

This study was approved by the Institutional Human Ethics Committee for Student Research (Care IHEC-I), India, wide reference number IHEC-II/0162/22.

INFORMED CONSENT

Written informed consent was obtained from the participants before enrolling in the study.

REFERENCES

- Maiocchi SL, Morris JC, Rees MD, Thomas SR. Regulation of the nitric oxide oxidase activity of myeloperoxidase by pharmacological agents. *Biochem Pharmacol.* 2017;135:90-115. doi: 10.1016/j. bcp.2017.03.016
- 2. Kavousi M. Subclinical Measures of Atherosclerosis: Genetics and Cardiovascular Risk Prediction. 2013.
- Mottillo S, Filion KB, Genest J, et al. The metabolic syndrome and cardiovascular risk: a systematic review and meta-analysis. J Am Coll Cardiol. 2010;56(14):1113-1132. doi: 10.1016/j.jacc.2010.05.034
- 4. Toth PP. Subclinical atherosclerosis: what it is, what it means and what we can do about it. *Int J Clin Pract.* 2008;62(8):1246-1254. doi: 10.1111/j.1742-1241.2008.01804.x
- 5. Duclos TW. Function of CRP. Ann Med. 2000;32(4):274-278. doi: 10.3109/07853890009011772
- Kim-Mitsuyama S, Soejima H, Yasuda O, et al. Reduction in hsCRP levels is associated with decreased incidence of cardiovascular events in Japanese hypertensive women but not in men. *Sci Rep.* 2020;10(1):17040. doi: 10.1038/s41598-020-73905-4
- Calabroi P, Willerson JT, Yeh ET. Inflammatory cytokines stimulated C-reactive protein production by human coronary artery smooth muscle cells. *Circulation*. 2003;108(16):1930-1932. doi: 10.1161/01. CIR.0000096055.62724.C5
- Ridker PM, Rifai N, Pfeffer M, Sacks F, Lepage S, Braunwald E. Elevation of tumor necrosis factor-α and increased risk of recurrent coronary events after myocardial infarction. *Circulation*. 2000;101(18):2149-2153. doi: 10.1161/01.CIR.101.18.2149
- 9. Lam JC, Ip MS. Sleep & the metabolic syndrome. *Indian Journal of Medical Research*. 2010;131(2):206-17.
- Handelsman Y. Metabolic syndrome pathophysiology and clinical presentation. *Toxicologic pathology*. 2009;37(1):18-20.
- 11. Haverkate E, Thompson SG, Pyke SD, Gallimore JR.

Production of C-reactive protein and risk of coronary events in stable and unstable angina. *Lancet*. 1997;349(9050):462-466. doi: 10.1016/S0140-6736(96)07591-5

- Liuzzo G, Biasucci LM, Gallimore JR, et al. The prognostic value of C-reactive protein and serum amyloid a protein in severe unstable angina. N Engl J Med. 1994;331(7):417-424. doi: 10.1056/ NEJM199408183310701
- de Winter RJ, Koch KT, van Straalen JP, et al. C-reactive protein and coronary events following percutaneous coronary angioplasty. *Am J Med.* 2003;115(2):85-90. doi: 10.1016/S0002-9343(03)00238-9
- 14. Morrow DA, de Lemos JA, Sabatine MS, et al. Clinical relevance of C-reactive protein during follow-up of patients with acute coronary syndromes in the Aggrastat-to-Zocor Trial. *Circulation.* 2006;114(4):281-288. doi: 10.1161/CIRCULATIONAHA.106.628909
- 15. Ridker PM. Clinical application of C-reactive protein for cardiovascular disease detection and prevention. *Circulation*. 2003;107(3):363-369. doi: 10.1161/01. CIR.0000053730.47739.3C
- Pai JK, Pischon T, Ma J, et al. Inflammatory markers and the risk of coronary heart disease in men and women. *N Engl J Med*. 2004;351(25):2599-610. doi: 10.1056/ NEJMoa040967
- 17. Tsimikas S, Willerson JT, Ridker PM. C-reactive protein and other emerging blood biomarkers to optimize risk stratification of vulnerable patients. *J Am Coll Cardiol.* 2006;47(8 suppl):C19-31. doi: 10.1016/j. jacc.2005.10.066
- Stewart LK, Earnest CP, Blair SN, Church TS. Effects of different doses of physical activity on C-reactive protein among women. *Med Sci Sports Exerc.* 2010;42(4):701-707. doi: 10.1249/MSS.0b013e3181c03a2b
- Ridker PM, Cushman M, Stampfer MJ, Tracy RP, Hennekens CH. Inflammation, aspirin, and the risk of cardiovascular disease in apparently healthy men. *N Engl J Med*. 1997;336(14):973-979. doi: 10.1056/ NEJM199704033361401
- Sattar N, Gaw A, Scherbakova O, et al. Metabolic syndrome with and without C-reactive protein as a predictor of coronary heart disease and diabetes in the West of Scotland Coronary Prevention Study. *Circulation*. 2003;108(4):414-449. doi: 10.1161/01. CIR.0000080897.52664.94
- Koenig W, Khuseyinova N. Lipoprotein-associated and secretory phospholipase A 2 in cardiovascular disease: the epidemiological evidence. *Cardiovasc Drugs Ther.* 2009;23(1):85-92. doi: 10.1007/s10557-008-6135-6
- Reilly MP, Wolfe ML, Localio AR, Rader DJ. C-reactive protein and coronary artery calcification: The Study of Inherited Risk of Coronary Atherosclerosis (SIRCA). *Arterioscler Thromb Vasc Biol.* 2003;23(10):1851-1856. doi: 10.1161/01.ATV.0000092327.60858.4A
- Hanley AJ, Festa A, D'Agostino Jr RB, et al. Metabolic and inflammation variable clusters and prediction of type 2 diabetes: factor analysis using directly measured insulin sensitivity. *Diabetes*. 2004;53(7):1773-81. doi: 10.2337/diabetes.53.7.1773
- 24. Brennan ML, Hazen SL. Emerging role of myeloperoxidase and oxidant stress markers in

cardiovascular risk assessment. *Curr Opin Lipidol.* 2003;14(4):353-359. doi: 10.1097/00041433-200308000-00003

- Hazen SL, Heinecke JW. 3-Chlorotyrosine, a specific marker of myeloperoxidase-catalyzed oxidation, is markedly elevated in low density lipoprotein isolated from human atherosclerotic intima. J Clin Invest. 1997;99(9):2075-2081. doi: 10.1172/JCl119379
- Daugherty A, Dunn JL, Rateri DL, Heinecke JW. Myeloperoxidase, a catalyst for lipoprotein oxidation, is expressed in human atherosclerotic lesions. J Clin Invest. 1994;94(1):437-444. doi: 10.1172/JCI117342
- Bergt C, Pennathur S, Fu X,et al. The myeloperoxidase product hypochlorous acid oxidizes HDL in the human artery wall and impairs ABCA1-dependent cholesterol transport. *Proc Natl Acad Sci.* 2004;101(35):13032-13037. doi: 10.1073/pnas.0405292101
- Pennathur S, Bergt C, Shao B, et al. Human atherosclerotic intima and blood of patients with established coronary artery disease contain high density lipoprotein damaged by reactive nitrogen species. J Biol Chem. 2004;279(41):42977-42983. doi: 10.1074/jbc.M406762200
- Zheng L, Nukuna B, Brennan ML, et al. Apolipoprotein Al is a selective target for myeloperoxidase-catalyzed oxidation and functional impairment in subjects with cardiovascular disease. J Clin Invest. 2004;114(4):529-541. doi: 10.1172/JCl200421109
- Sugiyama S, Okada Y, Sukhova GK, Virmani R, Heinecke JW, Libby P. Macrophage myeloperoxidase regulation by granulocyte macrophage colony-stimulating factor in human atherosclerosis and implications in acute coronary syndromes. *Am J Pathol.* 2001;158(3):879-891. doi: 10.1016/S0002-9440(10)64036-9
- Sugiyama S, Kugiyama K, Aikawa M, Nakamura S, Ogawa H, Libby P. Hypochlorous acid, a macrophage product, induces endothelial apoptosis and tissue factor expression: involvement of myeloperoxidase-mediated oxidant in plaque erosion and thrombogenesis. *Arterioscler Thromb Vasc Biol.* 2004;24(7):1309-14. doi: 10.1161/01.ATV.0000131784.50633.4f
- 32. Naruko T, Ueda M, Haze K, et al. Neutrophil infiltration of culprit lesions in acute coronary syndromes. *Circulation*. 2002;106(23):2894-900. doi: 10.1161/01. CIR.0000042674.89762.20
- Nikpoor B, Turecki G, Fournier C, Theroux P, Rouleau GA. A functional myeloperoxidase polymorphic variant is associated with coronary artery disease in French-Canadians. Am Heart J. 2001;142(2):336-339. doi: 10.1067/mhj.2001.116769
- Vita JA, Brennan ML, Gokce N, et al. Serum myeloperoxidase levels independently predict endothelial dysfunction in humans. *Circulation*. 2004;110(9):1134-1139. doi: 10.1161/01. CIR.0000140262.20831.8F
- Zhang R, Brennan ML, Fu X, et al. Association between myeloperoxidase levels and risk of coronary artery disease. JAMA. 2001;286(17):2136-2142. doi: 10.1001/jama.286.17.2136
- Baldus S, Heeschen C, Meinertz T, et al. Myeloperoxidase serum levels predict risk in patients with acute coronary syndromes. *Circulation*. 2003;108(12):1440-

1445. doi: 10.1161/01.CIR.0000090690.67322.51

- Brennan ML, Penn MS, Van Lente F, et al. Prognostic value of myeloperoxidase in patients with chest pain. *N Engl J Med*. 2003;349(17):1595-1604. doi: 10.1056/ NEJMoa035003
- Shishehbor MH, Brennan ML, Aviles RJ, et al. Statins promote potent systemic antioxidant effects through specific inflammatory pathways. *Circulation*. 2003;108(4):426-431. doi: 10.1161/01. CIR.0000080895.05158.8B
- Ronald JA. Imaging myeloperoxidase activity in cardiovascular disease. *Curr Cardiovasc Imaging Rep.* 2011;4:24-31. doi: 10.1007/s12410-010-9056-2
- Rosenson RS, Brewer Jr HB, Ansell BJ, et al. Dysfunctional HDL and atherosclerotic cardiovascular disease. Nat Rev Cardiol. 2016;13(1):48-60. doi: 10.1038/nrcardio.2015.124
- Hafiane A, Genest J. High density lipoproteins: measurement techniques and potential biomarkers of cardiovascular risk. *BBA Clinical*. 2015;3:175-188. doi: 10.1016/j.bbacli.2015.01.005
- Toprak A, Kandavar R, Toprak D, et al. C-reactive protein is an independent predictor for carotid artery intima-media thickness progression in asymptomatic younger adults (from the Bogalusa Heart Study). BMC Cardiovasc Disord. 2011;11(1):1-8. doi: 10.1186/1471-2261-11-78
- Mori T, Sasaki J, Kawaguchi H, et al. Serum glycoproteins and severity of coronary atherosclerosis. *Am Heart J.* 1995;129(2):234-238. doi: 10.1016/0002-8703(95)90003-9
- Iana A, Sirbu E. Linking myeloperoxidase with subclinical atherosclerosis in adults with metabolic syndrome. Wiener klinische Wochenschrift. 2020;132:150-154. doi: 10.1007/s00508-019-01602-y
- 45. Yang Q, Lin Q, Guo D, et al. Association of carotid intima media thickness with metabolic syndrome among low-income middle-aged and elderly Chinese: a population-based cross-sectional study. Front

Cardiovasc Med. 2021;8:669245. doi: 10.3389/ fcvm.2021.669245

- Zhao M, Caserta CA, Medeiros CC, et al. Metabolic syndrome, clustering of cardiovascular risk factors and high carotid intima-media thickness in children and adolescents. J Hypertens. 2020;38(4):618-624. doi: 10.1097/HJH.00000000002318
- Kwaifa IK, Bahari H, Yong YK, Noor SM. Endothelial dysfunction in obesity-induced inflammation: molecular mechanisms and clinical implications. *Biomolecules*. 2020;10(2):291. doi: 10.3390/ biom10020291
- Raele R, Lotufo PA, Bittencourt MS, et al. The association of waist-to-height ratio and other anthropometric measurements with subclinical atherosclerosis: Results from the Brazilian Longitudinal Study of Adult Health (ELSA-Brasil). Nutr Metab Cardiovasc Dis. 2020;30(11):1989-1998. doi: 10.1016/j. numecd.2020.05.025
- Koziarska-Rosciszewska M, Gluba-Brzozka A, Franczyk B, Rysz J. High-sensitivity C-reactive protein relationship with metabolic disorders and cardiovascular diseases risk factors. *Life*. 2021;11(8):742. doi: 10.3390/ life11080742
- Papageorgiou N, Briasoulis A, Androulakis E, Tousoulis D. Imaging subclinical atherosclerosis: where do we stand? *Curr Cardiol Rev.* 2017;13(1):47-55. doi: 10.21 74/1573403X12666160803095855
- Verma S, Wang CH, Li SH, et al. A self-fulfilling prophecy: C-reactive protein attenuates nitric oxide production and inhibits angiogenesis. *Circulation*. 2002;106(8):913-919. doi: 10.1161/01. CIR.0000029802.88087.5E
- 52. Fichtlscherer S, Rosenberger G, Walter DH, Breuer S, Dimmeler S, Zeiher AM. Elevated C-reactive protein levels and impaired endothelial vasoreactivity in patients with coronary artery disease. *Circulation*. 2000;102(9):1000-1006. doi: 10.1161/01. CIR.102.9.1000